

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase III Randomized, Open-Label Active Comparator-Controlled Multicenter Study to Evaluate Efficacy and Safety of Obinutuzumab in Patients With Primary Membranous Nephropathy
Short Title:	A Study Evaluating the Efficacy and Safety of Obinutuzumab in Participants With Primary Membranous Nephropathy
Protocol Number:	601107320058367332
NCT Number:	NCT04629248
Sponsor Name:	Roche
Investigational Product:	caffeine
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Percentage of Participants who Achieve a Complete Remission (CR) at Week 104
Secondary Obj.:	1. Percentage of Participants who Achieve an Overall Remission at Week 104 2. Percentage of Participants who Achieve CR at Week 76 3. Time to Treatment Failure, Meeting Escape Criteria, or Relapse after Complete or Partial Remission

2.2 Background & Rationale

This study will evaluate the efficacy, safety, pharmacodynamics, and pharmacokinetics (PK) of obinutuzumab compared with tacrolimus in participants with primary membranous nephropathy (pMN).

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: ACTIVE_NOT_RECRUITING
Target N:	142

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Diagnosis of primary membranous nephropathy (pMN) according to renal biopsy prior to or during screening
- * Screening urinary protein-to-creatinine ratio (UPCR) ≥ 5 g/g from 24-hour urine collection after best supportive care for ≥ 3 months prior to screening or screening UPCR ≥ 4 g/g after best supportive care for ≥ 6 months prior to screening
- * eGFR ≥ 40 mL/min/1.73m² or qualified endogenous creatinine clearance ≥ 40 mL/min/1.73m² based on 24-hour urine collection during screening
- * Other inclusion criteria may apply

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4.2 Exclusion Criteria

- * Participants with a secondary cause of MN
- * Pregnancy or breastfeeding
- * Evidence of $\geq 50\%$ reduction in proteinuria during the previous 6 months prior to randomization
- * Severe renal impairment, including the need for dialysis or renal replacement therapy
- * Type 1 or 2 diabetes mellitus
- * Receipt of an excluded therapy, including any anti-CD20 therapy less than 9 months prior to or during screening; or cyclophosphamide, tacrolimus, or cyclosporin less than 6 months prior to or during screening
- * Significant or uncontrolled medical disease which, in the investigator's opinion, would preclude participant participation
- * Known active infection of any kind or recent major episode of infection
- * Major surgery requiring hospitalization within the 4 weeks prior to screening
- * Current active alcohol or drug abuse or history of alcohol or drug abuse within 12 months prior to screening
- * Intolerance or contraindication to study therapies
- * Other exclusion criteria may apply

11. Study Identifiers & Governance

Primary ID: 601107320058367332

Registry Number: NCT04629248

Sponsor: Roche

Study Title: A Study Evaluating the Efficacy and Safety of Obinutuzumab in Participants With Primary Membranous Nephropathy

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____