

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase III, Randomized, Double-blind, Placebo-controlled Study Evaluating Atezolizumab and Bevacizumab, With or Without Tiragolumab, in Patients With Untreated Locally Advanced or Metastatic Hepatocellular Carcinoma
Short Title:	A Study Evaluating Atezolizumab and Bevacizumab, With or Without Tiragolumab, in Participants With Untreated Locally Advanced or Metastatic Hepatocellular Carcinoma (HCC) (IMbrave152)
Protocol Number:	1020926893565647374
NCT Number:	NCT05904886
Sponsor Name:	Roche
Investigational Product:	bevacizumab
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Investigator-assessed Progression-free Survival (PFS) According to Response Evaluation Criteria in Solid Tumors Version 1.1 (RECIST v1.1) 2. Overall Survival (OS)
Secondary Obj.:	1. Investigator-assessed Confirmed Objective Response Rate (ORR) According to RECIST v1.1 2. Investigator-assessed Duration of Response (DOR) According to RECIST v1.1 3. Investigator-assessed PFS Rate According to RECIST v1.1 at 6 and 12 Months

2.2 Background & Rationale

The purpose of this study is to assess the efficacy and safety of tiragolumab, an anti-TIGIT monoclonal antibody, when administered in combination with atezolizumab and bevacizumab as first-line treatment, in participants with unresectable, locally advanced or metastatic HCC.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: ACTIVE_NOT_RECRUITING
Target N:	687

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Locally advanced or metastatic and/or unresectable HCC with diagnosis confirmed by histology/cytology or clinically by American Association for the Study of Liver Diseases (AASLD) criteria in cirrhotic participants
- * Disease that is not amenable to curative surgical and/or locoregional therapies

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * No prior systemic treatment for locally advanced or metastatic and/or unresectable HCC-measurable disease according to RECIST v1.1
- * Eastern cooperative oncology group (ECOG) performance status of 0 or 1 within 7 days prior to randomization
- * Child-pugh Class A within 7 days prior to randomization
- * Adequate hematologic and end-organ function
- * Female participants of childbearing potential must be willing to avoid pregnancy within 5 months after the final dose of atezolizumab, within 6 months after the final dose of bevacizumab, and within 90 days after the final dose of tiragolumab/placebo
- * Male participants with a female partner of childbearing potential or pregnant female partner must remain abstinent or use a condom during the treatment period and for 6 months after the final dose of bevacizumab and for 90 days after the final dose of tiragolumab/placebo to avoid exposing the embryo.

4.2 Exclusion Criteria

- * Pregnancy or breastfeeding within 5 months after the final dose of atezolizumab, within 6 months after the final dose of bevacizumab, and within 90 days after the final dose of tiragolumab/placebo
- * Prior treatment with cluster of differentiation 137 (CD137) agonists or immune checkpoint blockade therapies
- * Treatment with investigational therapy within 28 days prior to initiation of study treatment
- * Treatment with locoregional therapy to liver within 28 days prior to initiation of study treatment, or non-recovery from side effects of any such procedure
- * Treatment with systemic immunostimulatory agents
- * Treatment with systemic immunosuppressive medication
- * Untreated or incompletely treated esophageal and/or gastric varices with bleeding or that are at high risk for bleeding
- * A prior bleeding event due to esophageal and/or gastric varices within 6 months prior to initiation of study treatment
- * Active or history of autoimmune disease or immune deficiency
- * History of idiopathic pulmonary fibrosis, organizing pneumonia, drug-induced pneumonitis, or idiopathic pneumonitis, or evidence of active pneumonitis on screening chest computed tomography (CT) scan
- * History of malignancy other than HCC within 5 years prior to screening, with the exception of malignancies with a negligible risk of metastasis or death
- * Mixed histology or other subtypes/variants of HCC, including, but not limited to, known liver adenocarcinoma, fibrolamellar HCC, sarcomatoid HCC, other rare HCC variant, or mixed cholangiocarcinoma and HCC
- * Co-infection with hepatitis B virus (HBV) and hepatitis C virus (HCV)
- * Acute epstein-barr virus (EBV) infection or known or suspected chronic active EBV infection
- * Symptomatic, untreated, or actively progressing central nervous system (CNS) metastases.

11. Study Identifiers & Governance

Primary ID:	1020926893565647374
Registry Number:	NCT05904886
Sponsor:	Roche
Study Title:	A Study Evaluating Atezolizumab and Bevacizumab, With or Without Tiragolumab, in Participants With Untreated Locally Advanced or Metastatic Hepatocellular Carcinoma (HCC) (IMbrave152)

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____