

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A 16-WEEK, MULTICENTER, INTERVENTIONAL, PHASE 3, RANDOMIZED, DOUBLE-BLIND, PLACEBO-CONTROLLED, PARALLEL GROUP STUDY TO INVESTIGATE EFFICACY AND SAFETY OF ABROCITINIB IN CHILDREN 6 TO LESS THAN 12 YEARS OF AGE WITH MODERATE-TO-SEVERE ATOPIC DERMATITIS
Short Title:	A Study of the Medicine Called Abrocitinib in Children 6 to Less Than 12 Years of Age With Moderate-to-Severe Eczema
Protocol Number:	1075556539970939144
NCT Number:	NCT06807268
Sponsor Name:	Pfizer
Investigational Product:	abrocitinib
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	- Response based on achieving Validated Investigator's Global Assessment (vIGA) score of clear (0) or almost clear (1) (on a 5-point scale) and a reduction from baseline of ≥2 points at Week 12: The difference in proportion of responders based on vIGA at Week 12 in patients with moderate-to-severe AD treated with abrocitinib versus placebo - Response based on achieving ≥75% improvement from baseline in the Eczema Area and Severity Index (EASI)-75 at Week 12: The difference in proportion of responders based on EASI-75 at Week 12 in patients with moderate-to-severe AD treated with abrocitinib versus placebo
Secondary Obj.:	- Change from Baseline (CFB) in the Worst Itch Numerical Rating Scale (WI-NRS) at Week 2 - Response based on achieving at least a 4-point improvement from baseline in the WI-NRS at Week 12 - Response based on achieving WI-NRS <2 at Week 12

2.2 Background & Rationale

This research study is being conducted to find out if the test medicine, abrocitinib, improves eczema and is safe for children 6 to <12 years of age who have moderate-to-severe eczema. Research study participants who meet the study criteria will be assigned by chance (like the flip of a coin) to receive either abrocitinib test medicine or placebo (pretend medicine that looks just like the test medicine) for 16 weeks. The study will last for about 24 weeks in total.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: RECRUITING
Target N:	150

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

Body weight ≥ 15 kg

Exclusion Criteria

Participants are excluded from the study if any of the following criteria apply:

Medical Conditions:

Any medical or psychiatric condition including any active suicidal ideation in the past year or suicidal behavior in the past 5 years or laboratory abnormality that may increase the risk of study participation or, in the investigator's judgment, make the participant inappropriate for the study.

If the participant has SDQ total score ≥ 17 , the investigator should exclude them or refer the child to a pediatric MHP to determine if it is safe to participate in the study. A copy or summary of the evaluation should be placed in the site source documents.

Have any of the following medical conditions:

* Infections:

- * Skin infections that require treatment with systemic antimicrobials within 2 weeks prior to Day 1 (Baseline) or have superficial skin infections within 1 week of Day 1.

- * History of systemic infection requiring hospitalization or parenteral antimicrobial therapy or as otherwise judged clinically significant by the investigator within 1 month prior to Day 1.

- * Have a history (single episode) of disseminated herpes zoster or disseminated herpes simplex, or a recurrent localized, dermatomal herpes zoster.

- * Infection with HIV, hepatitis B, and/or hepatitis C

- * Evidence of active TB or inadequately treated latent TB.

* Skin Conditions:

- * Including but not limited to psoriasis, seborrheic dermatitis or lupus on Day 1 that would interfere with evaluation of AD or response to treatment.

* Other Conditions:

- * Documented history of skeletal dysplasia.

- * Documented history of retinal detachment.

- * History of or conditions associated with thrombocytopenia, coagulopathy or platelet dysfunction.

- * Prior history of leukemia, lymphoma, sarcoma or any other malignancy.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Immunodeficiency disorder or a first-degree relative with a hereditary immunodeficiency.
- * Any other medical conditions that in the investigator's judgment make the participant inappropriate for the study.

Prior/Concomitant Therapy:

Prior treatment with a systemic JAK inhibitor for AD. Live attenuated vaccination within 6 weeks prior to Day 1 or require vaccination with live attenuated vaccines during treatment or within 6 weeks after the last dose of study intervention.

Concomitant use of strong inhibitors and inducers of CYP2C19 enzymes, strong inducers of CYP2C9 enzymes, P-gp substrates with narrow therapeutic index and sensitive CYP2C19 substrates is not allowed in the study.

Prior/Concurrent Clinical Study Experience:

Previous administration of an investigational drug within 30 days or 5 half lives, whichever is longer, of Day 1.

4.2 Exclusion Criteria

Not Provided

11. Study Identifiers & Governance

Primary ID:	1075556539970939144
Registry Number:	NCT06807268
Sponsor:	Pfizer
Study Title:	A Study of the Medicine Called Abrocitinib in Children 6 to Less Than 12 Years of Age With Moderate-to-Severe Eczema

15. Sign-off & Approval

Principal Investigator: _____	Date: _____
Clinical Operations Lead: _____	Date: _____
Lead Statistician: _____	Date: _____