

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Randomized, Double-blind, Double-dummy, Placebo-controlled, Multicenter, Phase 3 Study Assessing the Efficacy, Safety, and Tolerability of 2 Doses of Remibrutinib Over a 68-week Treatment Period in Adult Patients With Moderate to Severe Hidradenitis Suppurativa.
Short Title:	A Phase 3 Study to Assess Efficacy Safety and Tolerability of Remibrutinib in Adult Patients With Moderate to Severe Hidradenitis Suppurativa
Protocol Number:	1088022346333617332
NCT Number:	NCT06799000
Sponsor Name:	Novartis
Investigational Product:	ibrutinib
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Proportion of participants with Hidradenitis Suppurativa clinical response 50 (HiSCR50) at Week 16: HiSCR50 is defined as at least a 50% decrease in Abscess and Inflammatory Nodule (AN) count with no increase in the number of abscesses and in the number of draining tunnels/fistulae compared to baseline.
Secondary Obj.:	1. Proportion of participants with Abscesses and inflammatory nodules 50 (AN50) response at Week 16 2. Percentage change from baseline in International Hidradenitis Suppurativa Severity Score System (IHS4) at Week 16 3. Proportion of participants with HiSCR75 response at Week 16

2.2 Background & Rationale

The purpose of this study is to establish the efficacy, safety, and tolerability of remibrutinib (LOU064) Dose A and Dose B compared to placebo in participants with moderate to severe hidradenitis suppurativa (HS).

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: RECRUITING
Target N:	555

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- Male and female participants ? 18 years of age at the time of signing of the informed consent.
- Diagnosis of HS based on clinical history and physical examination for at least 6 months prior to the Baseline visit.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

3. Participants with moderate to severe HS defined as:

- * A total of at least 5 AN count (abscesses and/or inflammatory nodules) AND
- * Inflammatory lesions should affect at least 2 distinct anatomic areas (e.g., left and right axillae)

Key Exclusion Criteria:

1. Presence of more than 20 fistulae/tunnels (both draining and non-draining) in total at baseline.
2. Any active skin disease or conditions that may interfere with the assessment of HS.
3. Previous exposure to remibrutinib or other BTK inhibitors.
4. Use of other investigational drugs within 5 half-lives of enrollment, or within 30 days (for small molecules) prior to randomization, or until the pharmacodynamic effect has returned to baseline (for biologics), whichever is longer.
5. Significant bleeding risk or coagulation disorders.
6. History of gastrointestinal bleeding.
7. Requirement for anti-platelet (except for acetylsalicylic acid up to 100 mg/d or clopidogrel up to 75 mg/d) or anti-coagulant medication.
8. History or current hepatic disease.
9. Evidence of clinically significant cardiovascular, neurological, psychiatric, pulmonary, renal, hepatic, endocrine, metabolic, hematological disorders, gastrointestinal disease or immunodeficiency that, in the Investigator's opinion, would compromise the safety of the participant, interfere with the interpretation of the study results or otherwise preclude participation or protocol adherence of the participant.
10. History of hypersensitivity to any of the study drug constituents.
11. Known or suspected infectious disease that is active, chronic or recurrent which precludes the participant from participating in the trial as per investigator's assessment. These infectious diseases include and are not limited to opportunistic infections (e.g., tuberculosis, atypical mycobacterioses, listeriosis or aspergillosis) and/or known or suspected Human Immunodeficiency Virus (HIV) infection. Should it be required by local regulations and/or considered appropriate by the investigator, an HIV test can be performed to confirm eligibility.
12. History of live attenuated vaccine administration within 6 weeks prior to randomization or requirement to receive these vaccinations at any time while on study treatment.
13. Major surgery within 8 weeks prior to screening or planned surgery for the duration of the study.

Other protocol-defined inclusion/exclusion criteria may apply.

4.2 Exclusion Criteria

1. Presence of more than 20 fistulae/tunnels (both draining and non-draining) in total at baseline.
2. Any active skin disease or conditions that may interfere with the assessment of HS.
3. Previous exposure to remibrutinib or other BTK inhibitors.
4. Use of other investigational drugs within 5 half-lives of enrollment, or within 30 days (for small molecules) prior to randomization, or until the pharmacodynamic effect has returned to baseline (for biologics), whichever is longer.
5. Significant bleeding risk or coagulation disorders.
6. History of gastrointestinal bleeding.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

7. Requirement for anti-platelet (except for acetylsalicylic acid up to 100 mg/d or clopidogrel up to 75 mg/d) or anti-coagulant medication.
8. History or current hepatic disease.
9. Evidence of clinically significant cardiovascular, neurological, psychiatric, pulmonary, renal, hepatic, endocrine, metabolic, hematological disorders, gastrointestinal disease or immunodeficiency that, in the Investigator's opinion, would compromise the safety of the participant, interfere with the interpretation of the study results or otherwise preclude participation or protocol adherence of the participant.
10. History of hypersensitivity to any of the study drug constituents.
11. Known or suspected infectious disease that is active, chronic or recurrent which precludes the participant from participating in the trial as per investigator's assessment. These infectious diseases include and are not limited to opportunistic infections (e.g., tuberculosis, atypical mycobacterioses, listeriosis or aspergillosis) and/or known or suspected Human Immunodeficiency Virus (HIV) infection. Should it be required by local regulations and/or considered appropriate by the investigator, an HIV test can be performed to confirm eligibility.
12. History of live attenuated vaccine administration within 6 weeks prior to randomization or requirement to receive these vaccinations at any time while on study treatment.
13. Major surgery within 8 weeks prior to screening or planned surgery for the duration of the study.

Other protocol-defined inclusion/exclusion criteria may apply.

11. Study Identifiers & Governance

Primary ID:	1088022346333617332
Registry Number:	NCT06799000
Sponsor:	Novartis
Study Title:	A Phase 3 Study to Assess Efficacy Safety and Tolerability of Remibrutinib in Adult Patients With Moderate to Severe Hidradenitis Suppurativa

15. Sign-off & Approval

Principal Investigator: _____	Date: _____
Clinical Operations Lead: _____	Date: _____
Lead Statistician: _____	Date: _____