

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	An Open-label, Multicenter, Phase 3 Randomized, Active-Comparator-Controlled Clinical Study of Pembrolizumab (MK-3475) in Combination With Sacituzumab Govitecan Versus MK-3475 Monotherapy as First-line Treatment in Participants With PD L1 TPS Greater Than or Equal to 50% Metastatic Non-small Cell Lung Cancer (KEYNOTE D46/EVOKE-03)
Short Title:	Study of Pembrolizumab (MK-3475) Monotherapy Versus Sacituzumab Govitecan in Combination With Pembrolizumab for Participants With Metastatic Non-small Cell Lung Cancer (NSCLC) With Programmed Cell Death Ligand 1 (PD-L1) Tumor Proportion Score (TPS) ≥50% (MK-3475-D46)
Protocol Number:	282539189255488180
NCT Number:	NCT05609968
Sponsor Name:	Merck
Investigational Product:	pembrolizumab
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Progression-Free Survival (PFS): PFS is defined as the time from randomization to the first documented progressive disease (PD) or death due to any cause, whichever occurs first as per the Response Criteria in Solid Tumors Version 1.1 (RECIST 1.1). PD is defined as ≥20% increase in the sum of diameters of target lesions. In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of ≥5 mm. The appearance of one or more new lesions is also considered PD. RECIST 1.1 is modified to follow a maximum of 10 target lesions and a maximum of 5 target lesions per organ. PFS as assessed by blinded independent central review (BICR) will be presented. 2. Overall Survival (OS): OS is defined as the time from randomization to death due to any cause.
Secondary Obj.:	1. Objective Response (OR) 2. Duration of Response (DOR) 3. Change from Baseline in the Global Health Status/Quality of Life (Items 29 and 30) Combined Score on the European Organization for Research and Treatment of Cancer Quality of Life Questionnaire - Core 30 (EORTC QLQ-C30)

2.2 Background & Rationale

The purpose of this study is to compare pembrolizumab (MK-3475) in combination with sacituzumab govitecan with pembrolizumab alone with respect to progression-free survival (PFS) and overall survival (OS) per Response Evaluation Criteria in Solid Tumors Version 1.1 (RECIST 1.1) as assessed by blinded independent central review (BICR) among

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

adults with metastatic non-small cell lung cancer (NSCLC) with programmed cell death ligand 1 (PD-L1) tumor proportion score (TPS) $\geq 50\%$).

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE3, Status: RECRUITING

Target N: 614

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

The main inclusion criteria include but are not limited to the following:

- * Has a histologically or cytologically confirmed diagnosis of metastatic non-small cell lung cancer (NSCLC)
- * Has confirmation that epidermal growth factor receptor (EGFR), anaplastic lymphoma kinase 1 (ALK-1), or ROS proto-oncogene 1 (ROS-1)-directed therapy is not indicated as primary therapy
- * Has provided tumor tissue that demonstrates PD-L1 tumor proportion score (TPS) $\geq 50\%$ of tumor cells as assessed by immunohistochemistry (IHC) at a central laboratory
- * Has a life expectancy of at least 3 months

4.2 Exclusion Criteria

The main exclusion criteria include but are not limited to the following:

- * Has history of a second malignancy, unless potentially curative treatment has been completed with no evidence of malignancy for 3 years
- * Has received prior systemic chemotherapy or other targeted or biological antineoplastic therapy for their metastatic NSCLC
- * Has previously received treatment with Topoisomerase 1 inhibitors or Trop-2 targeted therapy
- * Has received prior therapy with an anti-programmed cell death 1 protein (anti-PD-1), anti-programmed cell death ligand 1 (anti-PD-L1), or anti anti- programmed cell death ligand 2 (PD-L2) agent or with an agent directed to another stimulatory or coinhibitory T-cell receptor
- * Has received prior radiotherapy within 2 weeks of start of study intervention or has radiation-related toxicities requiring corticosteroids
- * Has received radiation therapy to the lung that is ≥ 30 Gray (Gy) within 6 months of the first dose of study intervention
- * Has received a live or live-attenuated vaccine within 30 days before the first dose of study intervention
- * Has received an investigational agent or has used an investigational device within 4 weeks before study intervention administration
- * Has cardiac disease
 - * Myocardial infarction or unstable angina pectoris within 6 months of enrollment
 - * History of serious ventricular arrhythmia, high-grade atrioventricular block, or other cardiac arrhythmias requiring antiarrhythmic medications; history of QT interval prolongation

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * New York Heart Association (NYHA) Class III or greater congestive heart failure or left ventricular ejection fraction of $\leq 40\%$
- * Has active chronic inflammatory bowel disease
- * Has diagnosis of immunodeficiency or is receiving chronic systemic steroid therapy or any other form of immunosuppressive therapy within 7 days prior the first dose of study medication
- * Has known active central nervous system (CNS) metastases and/or carcinomatous meningitis
- * Has severe hypersensitivity (?Grade 3) to pembrolizumab or sacituzumab govitecan and/or any of their excipients
- * Has active autoimmune disease that has required systemic treatment in past 2 years except replacement therapy
- * History of (noninfectious) pneumonitis/interstitial lung disease that required steroids or has current pneumonitis/interstitial lung disease
- * Has active infection requiring systemic therapy
- * Has history of human immunodeficiency virus (HIV) infection
- * History of hepatitis B or known active hepatitis C virus infection
- * Has history or current evidence of any condition, therapy, laboratory abnormality, or other circumstance that might confound the results of the study or interfere with the participant's participation for the full duration of the study, such that it is not in the best interest of the participant to participate, in the opinion of the treating investigator
- * Have not adequately recovered from major surgery or have ongoing surgical complications

11. Study Identifiers & Governance

Primary ID: 282539189255488180

Registry Number: NCT05609968

Sponsor: Merck

Study Title: Study of Pembrolizumab (MK-3475) Monotherapy Versus Sacituzumab Govitecan in Combination With Pembrolizumab for Participants With Metastatic Non-small Cell Lung Cancer (NSCLC) With Programmed Cell Death Ligand 1 (PD-L1) Tumor Proportion Score (TPS) $\geq 50\%$ (MK-3475-D46)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____