

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A MULTICENTER, RANDOMIZED, DOUBLE-BLIND PHASE 3 STUDY OF PALBOCICLIB (ORAL CDK 4/6 INHIBITOR) PLUS LETROZOLE VERSUS PLACEBO PLUS LETROZOLE FOR THE TREATMENT OF PREVIOUSLY UNTREATED ASIAN POSTMENOPAUSAL WOMEN WITH ER (+), HER2 (-) ADVANCED BREAST CANCER
Short Title:	A Study Of Palbociclib (PD-0332991) + Letrozole VS. Placebo+ Letrozole For 1st Line Treatment Of Asian Postmenopausal Women With ER+/HER2- Advanced Breast Cancer [PALOMA-4]
Protocol Number:	315791112436234872
NCT Number:	NCT02297438
Sponsor Name:	Pfizer
Investigational Product:	letrozole
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Progression-Free Survival (PFS) Based on Investigator's Assessment: PFS was based on Kaplan-Meier estimates. PFS was defined as the time from the date of randomization to the date of the first documentation of objective progression of disease (PD) or death due to any cause in the absence of documented PD, whichever occurred first. PD is defined using Response Evaluation Criteria In Solid Tumors Criteria (RECIST v1.1) as a 20% increase in the sum of of diameters of target measurable lesions above the smallest sum observed (over baseline if no decrease in the sum is observed during therapy), with a minimum absolute increase of 5 mm. In this outcome measure, PFS was based on investigator's assessment.
Secondary Obj.:	1. Progression-Free Survival (PFS) Based on Blinded Independent Central Review (BICR) 2. Percentage of Participants Wiht Objective Response (OR) Based on Investigator Assessment 3. Percentage of Participants With Objective Response (OR) Based on Investigator Assessment (Participants With Measureable Disease at Baseline)

2.2 Background & Rationale

The study is designed to compare the clinical benefit following treatment with letrozole in combination with Palbociclib versus letrozole in combination with placebo in Asian postmenopausal women with ER(+)/HER2(-) advanced breast cancer who have not received prior systemic anti cancer therapies for their advanced/metastatic disease.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: COMPLETED
-------------------	--

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Target N: 340

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Adult Asian women with locoregionally recurrent or metastatic disease not amenable to curative therapy
- * Confirmed diagnosis of ER positive breast cancer
- * No prior systemic anti-cancer therapy for advanced ER+ disease
- * Postmenopausal women
- * Measurable disease as per Response Evaluation Criterion in Solid Tumors [RECIST] or bone-only disease
- * Eastern Cooperative Oncology Group [ECOG] 0-1
- * Adequate organ and marrow function
- * Patient must agree to provide tumor tissue

4.2 Exclusion Criteria

- * Confirmed diagnosis of HER2 positive disease
- * Patients with advanced, symptomatic, visceral spread that are at risk of life threatening complication in the short term
- * Known uncontrolled or symptomatic CNS metastases
- * Prior neoadjuvant or adjuvant treatment with a non steroidal aromatase inhibitor (ie, anastrozole or letrozole) with disease recurrence while on or within 12 months of completing treatment
- * Prior treatment with any CDK 4/6 inhibitor

11. Study Identifiers & Governance

Primary ID: 315791112436234872

Registry Number: NCT02297438

Sponsor: Pfizer

Study Title: A Study Of Palbociclib (PD-0332991) + Letrozole VS. Placebo+ Letrozole For 1st Line Treatment Of Asian Postmenopausal Women With ER+/HER2- Advanced Breast Cancer [PALOMA-4]

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____