

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 3 Randomized, Active-controlled, Open-label, Multicenter Study to Compare the Efficacy and Safety of MK-2870 Monotherapy Versus Treatment of Physician's Choice as Second-line Treatment for Participants With Recurrent or Metastatic Cervical Cancer (TroFuse-020/GOG-3101/ENGOT-cx20)
Short Title:	A Study to Compare Sacituzumab Tirumotecan (MK-2870) Monotherapy Versus Treatment of Physician's Choice as Second-line Treatment for Participants With Recurrent or Metastatic Cervical Cancer (MK-2870-020/TroFuse-020/Gog-3101/ENGOT-cx20)
Protocol Number:	330391244239558788
NCT Number:	NCT06459180
Sponsor Name:	Merck
Investigational Product:	gemcitabine
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Objective Response Rate (ORR) in Sacituzumab Tirumotecan Run-in: ORR is defined as the percentage of participants with Complete Response (CR: disappearance of all target lesions) or Partial Response (PR: at least a 30% decrease in the sum of diameters of target lesions) per Response Evaluation Criteria In Solid Tumors Version 1.1 (RECIST 1.1). The percentage of participants who experience CR or PR as assessed by Blinded Independent Central Review (BICR) will be presented. 2. Number of Participants Experiencing One or More Adverse Events (AEs) in Sacituzumab Tirumotecan Run-in: An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. An AE can therefore be any unfavorable and unintended sign (including an abnormal laboratory finding), symptom, or disease (new or exacerbated) temporally associated with the use of a study intervention. 3. Number of Participants Discontinuing Study Treatment Due to an AE in Sacituzumab Tirumotecan Run-in: An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. An AE can therefore be any unfavorable and unintended sign (including an abnormal laboratory finding), symptom, or disease (new or exacerbated) temporally associated with the use of a study intervention.
Secondary Obj.:	1. Progression-free Survival (PFS) in Phase 3 Portion 2. ORR in Phase 3 Portion 3. Duration of Response (DOR) in Phase 3 Portion

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

2.2 Background & Rationale

This study will have two phases: a sacituzumab tirumotecan safety run-in and a Phase 3 portion. The safety run-in phase will be used to evaluate the efficacy and safety of sacituzumab tirumotecan at the dose for evaluation in the Phase 3 portion. The purpose of this study is to compare the efficacy and safety of sacituzumab tirumotecan versus treatment of physician's choice as second-line treatment for participants with recurrent or metastatic cervical cancer in the Phase 3 portion.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE3, Status: RECRUITING

Target N: 686

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Has histologically-confirmed diagnosis of squamous cell carcinoma, adenosquamous carcinoma, or adenocarcinoma of the cervix
- * Must have recurrent or metastatic cervical cancer that has progressed on or after treatment with 1 prior line of systemic platinum doublet chemotherapy (with or without bevacizumab) AND must have received anti-PD-1/anti-PD-L1 therapy as part of prior cervical cancer regimens
- * Has measurable disease per Response Evaluation Criteria In Solid Tumors (RECIST) 1.1, as assessed by the investigator. Lesions situated in a previously irradiated area are considered measurable if progression has been shown in such lesions
- * Is assigned female sex at birth, at least 18 years of age at the time of providing the informed consent
- * Has ECOG performance status of 0 or 1 within 7 days before allocation for the Sacituzumab Tirumotecan Run-in or within 7 days before randomization for the Phase 3 portion
- * Has provided tumor tissue (most recent sample is preferred) from a core or excisional biopsy of a tumor lesion not previously irradiated
- * HIV-infected participants must have well controlled human immunodeficiency virus (HIV) on antiretroviral therapy (ART)
- * Participants who are hepatitis B surface antigen (HBsAg) positive are eligible if they have received hepatitis B virus (HBV) antiviral therapy for at least 4 weeks, and have undetectable HBV viral load prior to allocation (Sacituzumab Tirumotecan Run-in) or randomization (Phase 3 portion)
- * Participants with history of hepatitis C virus (HCV) infection are eligible if HCV viral load is undetectable at screening
- * Has adequate organ function

4.2 Exclusion Criteria

- * Has Grade ≥2 peripheral neuropathy
- * Has history of documented severe dry eye syndrome, severe Meibomian gland disease and/or blepharitis, or severe corneal disease that prevents/delays corneal healing
- * Has active inflammatory bowel disease requiring immunosuppressive medication or previous history of inflammatory bowel disease (eg, Crohn's disease, ulcerative colitis, or chronic diarrhea)

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Has uncontrolled, significant cardiovascular disease or cerebrovascular disease.
- * Received prior systemic anticancer therapy
- * Received prior radiotherapy within 2 weeks of start of study intervention, or has radiation-related toxicities, requiring corticosteroids
- * Received a live or live-attenuated vaccine within 30 days before the first dose of study intervention. Administration of killed vaccines is allowed
- * Has other histological subtypes of cervical cancer apart from squamous cell carcinoma, adenocarcinoma, or adenosquamous carcinoma (eg, carcinosarcoma), or has a diagnosis of nonepithelial cancer (eg, sarcoma, neuroendocrine tumors) of the cervix.
- * Known additional malignancy that is progressing or has required active treatment within the past 3 years
- * Known active central nervous system (CNS) metastases and/or carcinomatous meningitis
- * Active infection requiring systemic therapy
- * HIV-infected participants with a history of Kaposi's sarcoma and/or Multicentric Castleman's Disease
- * Concurrent active Hepatitis B and active Hepatitis C virus infection
- * Severe hypersensitivity (?Grade 3) to sacituzumab tirumotecan or treatment of physician's choice (TPC) and/or any of their excipients, or other biologic therapy
- * Participants who have not adequately recovered from major surgery or have ongoing surgical complications
- * Has a history of (noninfectious) pneumonitis/ILD that required steroids or has current pneumonitis/ILD

11. Study Identifiers & Governance

Primary ID: 330391244239558788

Registry Number: NCT06459180

Sponsor: Merck

Study Title: A Study to Compare Sacituzumab Tirumotecan (MK-2870) Monotherapy Versus Treatment of Physician's Choice as Second-line Treatment for Participants With Recurrent or Metastatic Cervical Cancer (MK-2870-020/TroFuse-020/Gog-3101/ENGOT-cx20)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____