

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase IIIb/IV, Multicenter, Randomized, Open-Label, Two-Arm Study to Investigate the Efficacy, Safety, and Durability of Faricimab Administered up to Every 24 Weeks in Patients With Neovascular Age-Related Macular Degeneration
Short Title:	A Study to Determine the Efficacy, Safety, and Durability of Faricimab in Participants With Neovascular Age-Related Macular Degeneration
Protocol Number:	396550255396827043
NCT Number:	NCT06795048
Sponsor Name:	Roche
Investigational Product:	faricimab
Phase:	PHASE4
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Change from Baseline in Best-Corrected Visual Acuity (BCVA) Score Averaged Over Weeks 44, 48, and 52
Secondary Obj.:	1. Change from Baseline in BCVA Score Averaged Over Weeks 92, 96, and 100 2. Change from Baseline in BCVA Score Over Time 3. Percentage of Participants Avoiding a Loss of ≥15 Letters in BCVA from Baseline Over Time

2.2 Background & Rationale

This study is a Phase IIIb/IV, multicenter, randomized, two-arm, open-label 100-week study to investigate the efficacy, safety, and durability of intravitreal 6-mg faricimab administered at up to 24-week intervals in patients with neovascular age-related macular degeneration (nAMD) that are treatment-naïve in the study eye.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE4, Status: ACTIVE_NOT_RECRUITING
Target N:	274

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Overtly healthy as determined by medical evaluation that includes medical history and physical examination
- * Agreement to adhere to the contraception requirements described in the protocol

Ocular Inclusion Criteria for Study Eye:

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Active treatment-naïve macular neovascularization (MNV) secondary to age-related macular degeneration (AMD), confirmed by the investigator based on the presence of intraretinal fluid (IRF) or subretinal fluid (SRF) affecting the central subfield on optical coherence tomography (OCT)
- * BCVA of 83 to 24 letters, inclusive (20/25 to 20/320 approximate Snellen equivalent, using the early treatment diabetic retinopathy study [ETDRS] protocol and addressed at the initial testing distance of 4 meters on Day 1)
- * Sufficiently clear ocular media and adequate pupillary dilation to allow acquisition of good quality retinal images to confirm diagnosis

4.2 Exclusion Criteria

Ocular Exclusion Criteria for Study Eye:

- * MNV due to causes other than nAMD, such as ocular histoplasmosis, trauma, pathological myopia, angioid streaks, choroidal rupture, or uveitis
- * Retinal pigment epithelial tear involving the macula on Day 1
- * Current vitreous hemorrhage on Day 1
- * Prior periocular pharmacological or IVT treatment (including faricimab, anti-vascular endothelial growth factor [VEGF], or complement inhibitor medication) for other retinal diseases

Ocular Exclusion Criteria for Fellow (Non-Study) Eye:

- * Participants who have a nonfunctioning fellow (non-study) eye, defined as either BCVA of hand motion or worse, or no physical presence of non-study eye (i.e., monocular), at both the screening and study Day 1 visits

Ocular Exclusion for Both Eyes:

- * History of idiopathic or autoimmune associated uveitis in either eye
- * Active ocular inflammation or suspected or active ocular or periocular infection in either eye on study Day 1

11. Study Identifiers & Governance

Primary ID:	396550255396827043
Registry Number:	NCT06795048
Sponsor:	Roche
Study Title:	A Study to Determine the Efficacy, Safety, and Durability of Faricimab in Participants With Neovascular Age-Related Macular Degeneration

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____