

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Randomized, Open-label, Multi-center Phase III Trial Comparing Tisagenlecleucel to Standard of Care in Adult Participants With Relapsed or Refractory Follicular Lymphoma (FL)
Short Title:	A Phase III Trial Comparing Tisagenlecleucel to Standard of Care (SoC) in Adult Participants With r/r Follicular Lymphoma
Protocol Number:	422866721853332151
NCT Number:	NCT05888493
Sponsor Name:	Novartis
Investigational Product:	prednisone
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Progression-free survival (PFS) determined by blinded independent review committee (BIRC): Progression free survival (PFS) based on Lugano response criteria, defined as time from randomization to the first of the following events to occur: * progressive disease (by BIRC) * death from any cause
Secondary Obj.:	1. Complete response rate (CRR) as assessed by BIRC (Key Secondary) 2. Overall response rate (ORR) by BIRC 3. Overall survival (OS)

2.2 Background & Rationale

This trial will compare tisagenlecleucel to standard of care in adult participants with relapsed or refractory (r/r) follicular lymphoma.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: RECRUITING
Target N:	108

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- Age ≥ 18 years at the date of signing the informed consent form.
- Follicular lymphoma grade 1, 2, or 3A confirmed histologically after latest relapse (local assessment).
- Relapsed or refractory disease after a second or later line of systemic therapy including an anti-CD20 antibody and an

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

alkylating agent.

4. Disease that is both active on Positron emission tomography (PET) scan (defined as a score of 4 or 5 on the Deauville 5-point scale) and measurable on Computed tomography (CT) scan.
5. ECOG performance status of 0, 1 or 2 at screening.
6. Adequate hematologic, renal, hepatic and pulmonary organ function at screening.
7. Must meet the institutional criteria to undergo leukapheresis (unless historical leukapheresis is available).
8. Must be eligible for treatment with the selected standard of care regimen.

4.2 Exclusion Criteria

1. Follicular lymphoma grade 3B or evidence of histologic transformation.
2. Prior treatment with anti-CD19 therapy, gene therapy, or adoptive T-cell therapy.
3. Active CNS involvement by malignancy.
4. Clinically significant active infection, presence of Human immunodeficiency virus (HIV) antibody or active hepatitis B or C.
5. Active neurological autoimmune or inflammatory disorders (e.g., Guillain-Barré syndrome).
6. Investigational medicinal product within the last 30 days or five half-lives (whichever is longer) prior to randomization.
7. Clinically significant cardiovascular conditions such as acute coronary syndrome, significant cardiac arrhythmias, heart failure or decreased LVEF.

Other protocol defined inclusion/exclusion criteria may apply

11. Study Identifiers & Governance

Primary ID:	422866721853332151
Registry Number:	NCT05888493
Sponsor:	Novartis
Study Title:	A Phase III Trial Comparing Tisagenlecleucel to Standard of Care (SoC) in Adult Participants With r/r Follicular Lymphoma

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____