

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Randomized, Open Label Study to Compare the Effect of Once Monthly Administration of Subcutaneous Mircera Versus Epoetin Alfa on Maintenance of Hemoglobin Levels, Safety and Tolerability in Dialysis Patients With Chronic Renal Anemia.
Short Title:	A Study of Subcutaneous Mircera for the Maintenance Treatment of Dialysis Patients With Chronic Renal Anemia.
Protocol Number:	510110912010584999
NCT Number:	NCT00560404
Sponsor Name:	Roche
Investigational Product:	epoetin alfa
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Percentage of Participants Maintaining Their Mean Hemoglobin Concentration Within Plus or Minus 1 Gram/Deciliter of Their Reference Hemoglobin and Between the Target Range During Efficacy Evaluation Period: The target hemoglobin (Hb) range was defined as Hb concentration (gram/deciliter [g/dL]) between 10.5 and 12.5 g/dL during the efficacy evaluation period (EEP). EEP was from Week 29 to Week 36.
Secondary Obj.:	1. Mean Change From Baseline in Hemoglobin Concentration Between Baseline and at the Efficacy Evaluation Period 2. Percentage of Participants Maintaining Individual Hemoglobin Concentration Within the Range of 10.5 - 12.5 Gram/Decilitre Throughout the Efficacy Evaluation Period 3. Mean Time Spent in Hemoglobin Range of 10.5 - 12.5 Gram/Decilitre During the Efficacy Evaluation Period

2.2 Background & Rationale

This 2 arm study will compare the efficacy and safety of monthly administration of subcutaneous Mircera versus epoetin alfa for the maintenance of hemoglobin levels in dialysis patients with chronic renal anemia. Patients currently receiving maintenance treatment with epoetin alfa will be randomized to receive either monthly injections of Mircera with a starting dose (120, 200 or 360 micrograms) derived from the dose of epoetin alfa they were receiving in the week preceding study start, or to continue on epoetin alfa treatment. The anticipated time on study treatment is 3-12 months, and the target sample size is 100-500 individuals.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: COMPLETED
Target N:	233

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * adult patients, ≥ 18 years of age;
- * chronic renal anemia;
- * regular hemodialysis with the same schedule of dialysis for ≥ 12 weeks;
- * maintenance therapy with subcutaneous epoetin alfa at the same administration interval for 4 weeks.

4.2 Exclusion Criteria

- * transfusion of red blood cells during previous 2 months;
- * poorly controlled hypertension requiring interruption of epoetin alfa in previous 6 months;
- * acute or chronic bleeding;
- * active malignant disease (except non-melanoma skin cancer).

11. Study Identifiers & Governance

Primary ID: 510110912010584999

Registry Number: NCT00560404

Sponsor: Roche

Study Title: A Study of Subcutaneous Mircera for the Maintenance Treatment of Dialysis Patients With Chronic Renal Anemia.

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____