

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase III, Randomized, Open-label, Active-controlled, Multicenter Study Evaluating the Safety, Pharmacokinetics, Pharmacodynamic and Efficacy of Crovalimab Versus Eculizumab in Patients With Paroxysmal Nocturnal Hemoglobinuria (PNH) Currently Treated With Complement Inhibitors
Short Title:	A Study Evaluating the Safety, Pharmacokinetics, and Efficacy of Crovalimab Versus Eculizumab in Participants With Paroxysmal Nocturnal Hemoglobinuria (PNH) Currently Treated With Complement Inhibitors
Protocol Number:	662341437294610788
NCT Number:	NCT04432584
Sponsor Name:	Roche
Investigational Product:	eculizumab
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Percentage of Participants With Adverse Events (AEs) and by Severity: Severity determined according to National Cancer Institute (NCI) Common Terminology Criteria for Adverse Events, Version 5 (CTCAE v5). 2. Percentage of Participants With Injection-site Reactions, Infusion-related Reactions, Hypersensitivity and Infections (including Meningococcal Meningitis) 3. Percentage of Participants With Adverse Events (AEs) Leading to Study Drug Discontinuation
Secondary Obj.:	1. Serum Concentrations of Crovalimab or Eculizumab Over Time 2. Serum Concentrations of Ravulizumab at the Time of Crovalimab Initiation 3. Percentage of Participants With Anti-crovalimab Antibodies

2.2 Background & Rationale

A study designed to evaluate the safety of crovalimab with eculizumab in participants with PNH currently treated with complement inhibitors. This study will enroll approximately 190 participants.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: ACTIVE_NOT_RECRUITING
Target N:	190

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Body weight $\geq$ 40 kg at screening (pediatric participants with body weight $<$ 40 kg)
- * Treated with eculizumab or ravulizumab for PNH for at least 3 months prior to Day 1
- * Lactate Dehydrogenase Levels $\geq$ 2x the upper limit of normal (ULN) at screening
- * Willingness and ability to comply with all study visits and procedures
- * Documented diagnosis of PNH, confirmed by high sensitivity flow cytometry
- * Vaccination against Neisseria meningitidis serotypes A, C, W, and Y $<$ 3 years prior to initiation of study treatment; or, if not previously done, vaccination administered no later than one week after the first drug administration
- * Women of childbearing potential: agreement to remain abstinent (refrain from heterosexual intercourse) or use contraception during the treatment period and for 10.5 months after the final dose of crovalimab or for 3 months after the final dose of eculizumab (or longer if required by the local product label)

4.2 Exclusion Criteria

- * History of allogeneic bone marrow transplantation
- * History of myelodysplastic syndrome with Revised International Prognostic Scoring System (IPSS-R) prognostic risk categories of intermediate, high and very high
- * Pregnant or breastfeeding, or intending to become pregnant during the study, within 10.5 months after the final dose of crovalimab, or 3 months after the final dose of eculizumab (or longer if required by the local product label)
- * Participation in another interventional treatment study with an investigational agent or use of any experimental therapy within 28 days of screening or within 5 half-lives of that investigational product, whichever was greater: participants enrolled in an eculizumab or ravulizumab interventional study are eligible provided they fulfill eligibility (e.g., are willing and able to comply with the study assessments) and stop their participation in current trial before randomisation/enrolment
- * Positive for Active Hepatitis B and C infection (HBV/HCV)
- * Concurrent disease, treatment, procedure, or surgery or abnormality in clinical laboratory tests that could interfere with the conduct of the study, may pose any additional risk for the participant, or would, in the opinion of the investigator, preclude the participant's safe participation in and completion of the study
- * History of or ongoing cryoglobulinemia at screening

11. Study Identifiers & Governance

Primary ID: 662341437294610788

Registry Number: NCT04432584

Sponsor: Roche

Study Title: A Study Evaluating the Safety, Pharmacokinetics, and Efficacy of Crovalimab Versus Eculizumab in Participants With Paroxysmal Nocturnal Hemoglobinuria (PNH) Currently Treated With Complement Inhibitors

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____