

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase III, Multicenter, Single Arm Study Evaluating the Efficacy, Safety, Pharmacokinetics, and Pharmacodynamics of Crovalimab in Patients With Paroxysmal Nocturnal Hemoglobinuria (PNH) Not Previously Treated With Complement Inhibition
Short Title:	A Study Evaluating the Efficacy, Safety, Pharmacokinetics and Pharmacodynamics of Crovalimab in Participants With Paroxysmal Nocturnal Hemoglobinuria (PNH) Not Previously Treated With Complement Inhibition
Protocol Number:	669863048493186011
NCT Number:	NCT04654468
Sponsor Name:	Roche
Investigational Product:	Crovalimab
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Mean Percentage of Participants With Hemolysis Control: A Generalized Estimating Equation (GEE) was used to estimate the population-average percentage of the participants with hemolysis control (measured by lactate dehydrogenase (LDH) $\geq 1.5 \times$ upper limit of normal (ULN)) from Week 5 to Week 25 taking account of the intra-patient and inter-patient correlation between LDH control statuses across visits. The dependent variable was the binary indicator for hemolysis control. Independent variables are categorical effects of visits, continuous baseline LDH. 2. Difference in Percentage of Participants With Transfusion Avoidance (TA) From Baseline Through Week 25 and Within 24 Weeks Prior to Screening: TA was defined as participants who were packed red blood cell (pRBC) transfusion-free and did not require transfusion per protocol-specified guidelines. TA within 24 weeks prior to screening was based on the pRBC transfusion history in the medical records. Reported in this outcome measure is the difference in the percentage of participants between "baseline through Week 25" and "within 24 weeks prior to screening". 95% Confidence Interval (CI) for the difference between the percentage of participants with transfusion avoidance between Pre-screening and Post-baseline is calculated using the Newcombe method. Screening= Day -28 to Day -1 and Baseline= Day 1.
Secondary Obj.:	1. Percentage of Participants With Breakthrough Hemolysis (BTH) 2. Percentage of Participants With Stabilized Hemoglobin 3. Change From Baseline in Fatigue in Adults Aged ≥ 18 Years

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

2.2 Background & Rationale

This study will enroll participants aged 12 years or older with a body weight $\geq$ 40 kilograms (kg) diagnosed with PNH who have not been previously treated with complement inhibitor therapy. Approximately 50 participants will be treated with Crovalimab for at least 24 weeks.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE3, Status: ACTIVE_NOT_RECRUITING

Target N: 51

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Body weight $\geq$ 40 kg at screening
- * Willingness and ability to comply with all study visits and procedures
- * Documented diagnosis of PNH, confirmed by high sensitivity flow cytometry
- * LDH Levels $\geq$ 2x the ULN at screening
- * Participants who have at least four transfusions during 12 months prior to screening (documented in the medical record)
- * Presence of one or more of the PNH-related signs or symptoms within 3 months of screening
- * Vaccination against Neisseria meningitidis serotypes A, C, W, and Y $\leq$ 3 years prior to initiation of study treatment (Day 1)
- * Vaccination against Haemophilus influenzae type B and Streptococcus pneumonia according to national vaccination recommendations
- * For participants receiving other therapies (e.g., immunosuppressants, corticosteroids): stable dose for $\geq$ 28 days prior to screening and up to the first drug administration
- * Adequate hepatic and renal function
- * Women of childbearing potential: agreement to remain abstinent (refrain from heterosexual intercourse) or use contraception during the treatment period and for 46 weeks (approximately 10.5 months) after the final dose of crovalimab
- * Platelet count $\geq$ 30,000 per cubic millimeter (mm^3) at screening
- * ANC $\geq$ 500/microlitres (μL) at screening

4.2 Exclusion Criteria

- * Current or previous treatment with a complement inhibitor
- * History of allogeneic bone marrow transplantation
- * History of Neisseria meningitidis infection within 6 months prior to screening and up to first drug administration
- * Known or suspected immune or hereditary complement deficiency
- * Known HIV infection with cluster of differentiation 4 (CD4) count $\leq$ 200 cells/ μL within 24 weeks prior to screening
- * Infection requiring hospitalization or treatment with IV antibiotics within 28 days prior to screening and up to the first drug administration, or oral antibiotics within 14 days prior to screening and up to the first drug administration
- * Active systemic bacterial, viral, or fungal infection within 14 days before first drug administration

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Presence of fever ($\geq 38^{\circ}\text{C}$) within 7 days before the first drug administration
- * Splenectomy ≤ 6 months before screening
- * History of malignancy within 5 years prior to screening and up to the first drug administration
- * Pregnant or intending to become pregnant during the study or within 46 weeks (10.5 months) after the final dose of study treatment
- * Participation in another interventional treatment study with an investigational agent or use of any experimental therapy within 28 days of screening or within 5 half-lives of that investigational product, whichever is greater

11. Study Identifiers & Governance

Primary ID: 669863048493186011

Registry Number: NCT04654468

Sponsor: Roche

Study Title: A Study Evaluating the Efficacy, Safety, Pharmacokinetics and Pharmacodynamics of Crovalimab in Participants With Paroxysmal Nocturnal Hemoglobinuria (PNH) Not Previously Treated With Complement Inhibition

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____