

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase III, Open-label, Multi-center, Randomized Study Comparing AAA817+ARPI Versus Standard of Care in Adult Participants With PSMA-positive Metastatic Castration Resistant Prostate Cancer
Short Title:	Study Comparing AAA817+ARPI Versus Standard of Care in Adult Participants With PSMA-positive mCRPC
Protocol Number:	868829634600770456
NCT Number:	NCT06855277
Sponsor Name:	Novartis
Investigational Product:	procyclidine
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Radiographic Progression Free Survival (rPFS): Time to radiographic disease progression or death due to any cause.
Secondary Obj.:	1. Overall Survival (OS) (Key Secondary Endpoint) 2. Radiographic Progression Free Survival by PSMA PET/CT and (rPSF-PET)(Key Secondary) 3. Progression Free Survival (PFS)

2.2 Background & Rationale

The purpose of this study is to determine whether ^{225}Ac -PSMA-617 (AAA817), given for up to 6 cycles at a dose of 10 Megabecquerel (MBq) +/- 10%, plus androgen receptor pathway inhibitor (ARPI), improves the radiographic progression free survival (rPFS) compared to investigator's choice of standard of care (SOC) (ARPI change or taxane-based chemotherapy) in adult participants with PSMA-positive metastatic castration resistant prostate cancer (mCRPC) treated with another ARPI as last treatment and who have not been exposed to a taxane-containing chemotherapy in the mCRPC setting nor have received any prior PSMA-targeting radioligand therapy.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: RECRUITING
Target N:	605

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Signed informed consent must be obtained prior to participation in the study.
- * Participants must be adults ? 18 years of age.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Participants must have an ECOG performance status of 0 to 2.
- * Participants must have histological, and/or cytological confirmation of adenocarcinoma of the prostate. Participants with mixed histology (neuroendocrine) are not eligible.
- * Participants who have received taxane-based chemotherapy in mHSPC setting are eligible if they are deemed appropriate for chemotherapy or ARPI change as the next line of therapy in the opinion of the Investigator. Note: Participants who have received taxane-based chemotherapy for mCRPC are excluded.
- * Participants must not have received taxane-based chemotherapy in mCRPC setting (allowed in mHSPC setting).
- * Participants must have PSMA-PET positive disease using a PSMA imaging agent that is approved as per protocol.
- * Participant must have been diagnosed with mCRPC with documented progressive disease while on treatment with ARPI in mHSPC or earlier setting as their last treatment (and did not progress on more than one ARPI).

Key Exclusion Criteria:

- * Previous treatment with any approved or investigational RLT, approved or investigational radioisotopes
- * Previous treatment with any conventional external beam radiotherapy including hemi-body radiation within 6 weeks of randomization (within 2 weeks for radiotherapy of localized metastases).
- * Participants with known or suspected deleterious germline or somatic homologous recombination repair (HRR) gene-mutated metastatic castration-resistant prostate cancer.
- * Any approved or investigational agents/systemic anti-cancer therapy (e.g. other chemotherapy, investigational therapy, immunotherapy or biological therapy including monoclonal antibodies) within 28 days (or 5 times the half-life of that therapy whichever is longer) of the anticipated day C1D1.
- * Diagnosed with other active malignancies that are expected to alter life expectancy or may interfere with disease assessment.

Other protocol-defined inclusion/exclusion criteria may apply.

4.2 Exclusion Criteria

- * Previous treatment with any approved or investigational RLT, approved or investigational radioisotopes
- * Previous treatment with any conventional external beam radiotherapy including hemi-body radiation within 6 weeks of randomization (within 2 weeks for radiotherapy of localized metastases).
- * Participants with known or suspected deleterious germline or somatic homologous recombination repair (HRR) gene-mutated metastatic castration-resistant prostate cancer.
- * Any approved or investigational agents/systemic anti-cancer therapy (e.g. other chemotherapy, investigational therapy, immunotherapy or biological therapy including monoclonal antibodies) within 28 days (or 5 times the half-life of that therapy whichever is longer) of the anticipated day C1D1.
- * Diagnosed with other active malignancies that are expected to alter life expectancy or may interfere with disease assessment.

Other protocol-defined inclusion/exclusion criteria may apply.

11. Study Identifiers & Governance

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Primary ID: 868829634600770456
Registry Number: NCT06855277
Sponsor: Novartis
Study Title: Study Comparing AAA817+ARPI Versus Standard of Care in Adult Participants With PSMA-positive mCRPC

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____