

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase III, Randomized, Double-blind, Placebo-controlled, Multicenter Basket Study to Evaluate the Efficacy, Safety, Pharmacokinetics, and Pharmacodynamics of Satralizumab in Patients With Anti-N-methyl-D-aspartic Acid Receptor (NMDAR) or Anti-leucine-rich Glioma-inactivated 1 (LGI1) Encephalitis
Short Title:	A Study to Evaluate the Efficacy, Safety, Pharmacokinetics (PK), and Pharmacodynamics (PD) of Satralizumab in Participants With Anti-N-methyl-D-aspartic Acid Receptor (NMDAR) or Anti-leucine-rich Glioma-inactivated 1 (LGI1) Encephalitis
Protocol Number:	889225379838246316
NCT Number:	NCT05503264
Sponsor Name:	Roche
Investigational Product:	satralizumab
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Part 1: Proportion of Participants in NMDAR AIE Cohort With Modified Rankin Scale (mRS) Score Improvement ? 1 From Baseline and no Use of Rescue Therapy at Week 24 2. Part 1: Proportion of Participants in LGI1 AIE Cohort With mRS Score Improvement ? 1 From Baseline and no Use of Rescue Therapy at Week 52 3. Part 2: Percentage of Participants With Adverse Events (AEs)
Secondary Obj.:	1. Part 1 (NMDAR AIE Cohort and LGI1 Cohort): Time to mRS Score Improvement ? 1 From Baseline Without Use of Rescue Therapy 2. Part 1 (NMDAR AIE Cohort and LGI1 AIE Cohort): Time to Rescue Therapy 3. Part 1 (NMDAR AIE Cohort and LGI1 AIE Cohort): Proportion of Participants With Sustained Seizure Cessation at Week 24

2.2 Background & Rationale

The purpose of this study is to assess the efficacy, safety, PK, and PD of satralizumab in participants with NMDAR and LGI1 encephalitis.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: RECRUITING
Target N:	152

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Reasonable exclusion of tumor or malignancy before baseline visit (randomization)
- * Onset of AIE symptoms ? 9 months before randomization
- * Meet the definition of "New Onset" or "Incomplete Responder" AIE
- * For women of childbearing potential: agreement to remain abstinent (refrain from heterosexual intercourse) or use adequate contraception during the treatment period and for at least 3 months after the final dose of satralizumab or placebo
- * For participants enrolled in the extended China enrollment phase at China's sites: participants who are current residents of mainland China, Hong Kong, or Taiwan, and of Chinese ancestry

NMDAR AIE Cohort:

- * Age ? 12 years
- * Diagnosis of probable or definite NMDAR encephalitis

LGI1 AIE Cohort

- * Age ? 18 years
- * Diagnosis of LGI1 encephalitis

4.2 Exclusion Criteria

- * Any untreated teratoma or thymoma at baseline visit (randomization)
- * History of carcinoma or malignancy, unless deemed cured by adequate treatment with no evidence of recurrence for ? 5 years before screening
- * For participants with NMDAR AIE, history of negative anti-NMDAR antibody in cerebrospinal fluid (CSF) using a cell-based assay within 9 months of symptom onset
- * Historically known positivity to an intracellular antigen with high cancer association or glutamate decarboxylase 65 (GAD-65)
- * Historically known positivity to any cell surface neuronal antibodies other than NMDAR and LGI1, in the absence of NMDAR and LGI1 antibody positivity
- * Confirmed paraneoplastic encephalitis
- * Confirmed central or peripheral nervous system demyelinating disease
- * Alternative causes of associated symptoms
- * History of herpes simplex virus encephalitis in the previous 24 weeks
- * Any previous/concurrent treatment with interleukin-6 (IL-6) inhibitory therapy (e.g., tocilizumab), alemtuzumab, total body irradiation, or bone marrow transplantation
- * Any previous treatment with anti-cluster of differentiation 19 antibody (CD19 antibody), complement inhibitors, neonatal Fc receptor antagonists, anti-B-lymphocyte stimulator monoclonal antibody
- * Any previous treatment with T-cell depleting therapies, cladribine, or mitoxantrone
- * Treatment with oral cyclophosphamide within 1 year prior to baseline
- * Treatment with any investigational drug (including bortezomib) within 24 weeks prior to screening
- * Concurrent use of more than one immunosuppressive therapy (IST) as background therapy

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Contraindication to all of the following rescue treatments: rituximab, intravenous immunoglobulin (IVIG), high-dose corticosteroids, or intravenous (IV) cyclophosphamide
- * Any surgical procedure, except laparoscopic surgery or minor surgeries within 4 weeks prior to baseline, excluding surgery for thymoma or teratoma removal
- * Planned surgical procedure during the study
- * Evidence of progressive multifocal leukoencephalopathy
- * Evidence of serious uncontrolled concomitant diseases
- * Congenital or acquired immunodeficiency, including human immunodeficiency virus (HIV) infection
- * Active or presence of recurrent bacterial, viral, fungal, mycobacterial infection, or other infection
- * Infection requiring hospitalization or treatment with IV anti-infective agents within 4 weeks prior to baseline visit
- * Positive hepatitis B (HBV) and hepatitis C (HCV) test at screening
- * Evidence of latent or active tuberculosis (TB)
- * History of drug or alcohol abuse within 1 year prior to baseline
- * History of diverticulitis or concurrent severe gastrointestinal (GI) disorders that, in the investigator's opinion, may lead to increased risk of complications such as GI perforation
- * Receipt of live or live-attenuated vaccine within 6 weeks prior to baseline visit
- * History of blood donation (1 unit or more), plasma donation or platelet donation within 90 days prior to screening
- * History of severe allergic reaction to a biologic agent
- * History of suicide attempt within 3 years prior to screening except if this is clearly associated with and occurs during the acute phase of LGI-1 or NMDAR encephalitis
- * Any serious medical condition or abnormality in clinical laboratory tests that, in the investigator's judgment, precludes safe participation in and completion of the study
- * Pregnant or breastfeeding, or intending to become pregnant during the study or within 3 months after the final dose of study drug

11. Study Identifiers & Governance

Primary ID: 889225379838246316

Registry Number: NCT05503264

Sponsor: Roche

Study Title: A Study to Evaluate the Efficacy, Safety, Pharmacokinetics (PK), and Pharmacodynamics (PD) of Satralizumab in Participants With Anti-N-methyl-D-aspartic Acid Receptor (NMDAR) or Anti-leucine-rich Glioma-inactivated 1 (LGI1) Encephalitis

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____