

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 3, Randomized, Open-label Study to Evaluate the Efficacy and Safety of Sac-TMT (Sacituzumab Tirumotecan, MK-2870) Followed by Carboplatin/Paclitaxel vs Chemotherapy, Both in Combination With Pembrolizumab as Neoadjuvant Therapy for High-Risk, Early-Stage, Triple-Negative Breast Cancer or Hormone Receptor-low Positive/Human Epidermal Growth Factor Receptor-2 Negative Breast Cancer
Short Title:	A Clinical Study of Sacituzumab Tirumotecan (Sac-TMT, MK-2870) in People With Breast Cancer (MK-2870-032)
Protocol Number:	968994899850358315
NCT Number:	NCT06966700
Sponsor Name:	Merck
Investigational Product:	cyclophosphamide
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Percentage of Participants with Pathological Complete Response (pCR) at the Time of Definitive Surgery: pCR (ypT0/Tis ypN0) is defined as the absence of residual invasive cancer on hematoxylin and eosin evaluation of the complete resected breast specimen and all sampled regional lymph nodes after completion of neoadjuvant systemic therapy per current American Joint Committee on Cancer (AJCC) staging criteria assessed by the local pathologist at the time of definitive surgery. 2. Event-Free Survival (EFS): EFS is defined as the time from randomization to disease progression that precludes surgery, local or distant recurrence, or death due to any cause, whichever occurs first.
Secondary Obj.:	1. Overall Survival (OS) 2. Percentage of Participants with pCR with No Ductal Carcinoma in Situ (pCR-no DCIS) at the Time of Definitive Surgery 3. Percentage of Participants with pCR at the Time of Definitive Surgery (High-risk, early-stage, TNBC subset)

2.2 Background & Rationale

Researchers are looking for new ways to treat types of breast cancer that are both:

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: RECRUITING
Target N:	2400

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

The main inclusion criteria include but are not limited to the following:

* Has previously untreated high-risk, early-stage, non-metastatic (M0) breast cancer (BC), defined as any of the following combined primary tumor (T) and regional lymph node (N) staging per AJCC 8th edition criteria as assessed by the investigator based on radiological and/or clinical assessment:

- * cT1c, N1-N2
- * cT2, N0-N2
- * cT3, N0-N2
- * cT4a-d, N0-N2

* The participant must have a centrally confirmed diagnosis of BC that is triple-negative or HR-low+/HER2- (defined as estrogen receptor (ER)-low+ expression in 1% to 10% cells and HER2-), as by the most recent American Society of Clinical Oncology (ASCO)/College of American Pathologists (CAP) guidelines.

* Provides a core needle biopsy from the primary breast tumor at screening to the central laboratory.

* Has Eastern Cooperative Oncology Group (ECOG) performance status of 0 or 1 performed within 28 days before treatment randomization.

* Demonstrates adequate organ function.

4.2 Exclusion Criteria

The main exclusion criteria include but are not limited to the following:

* Metastatic (Stage IV) breast cancer or clinical node stage 3 (cN3) nodal involvement

* Has received any prior treatment, including radiation, systemic therapy, and/or definitive surgery for currently diagnosed breast cancer

* Has undergone excisional biopsy of the primary tumor, axillary lymph node dissection, and/or axillary sentinel lymph node biopsy prior to study treatment.

* Received prior systemic anticancer therapy including investigational agents within 4 weeks before randomization.

* Received prior therapy with an anti-PD-1, anti-PD-L1, or anti-PD-L2 agent, or with an agent directed to another stimulatory or coinhibitory T-cell receptor (eg, CTLA-4, OX-40, CD137).

* Received prior treatment with a TROP2-targeted antibody-drug conjugate (ADC).

* Received prior treatment with a topoisomerase I inhibitor-containing ADC.

* Received a live or live-attenuated vaccine within 30 days before the first dose of study intervention.

* Known additional malignancy that is progressing or has required active treatment within the past 5 years.

* Uncontrolled systemic disease.

* History of (noninfectious) pneumonitis/interstitial lung disease that required steroids or has current pneumonitis/interstitial lung disease.

11. Study Identifiers & Governance

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Primary ID: 968994899850358315
Registry Number: NCT06966700
Sponsor: Merck
Study Title: A Clinical Study of Sacituzumab Tirumotecan (Sac-TMT, MK-2870) in People With Breast Cancer (MK-2870-032)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____