

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	Adjuvant Immunotherapy With Anti-PD-1 Monoclonal Antibody Pembrolizumab (MK-3475) Versus Placebo After Complete Resection of High-risk Stage III Melanoma: A Randomized, Double- Blind Phase 3 Trial of the EORTC Melanoma Group
Short Title:	Study of Pembrolizumab (MK-3475) Versus Placebo After Complete Resection of High-Risk Stage III Melanoma (MK-3475-054/1325-MG/KEYNOTE-054)
Protocol Number:	1137004121105929006
NCT Number:	NCT02362594
Sponsor Name:	Merck
Investigational Product:	pembrolizumab
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Part 1: Percentage of Participants With Recurrence-Free Survival (RFS) At 6 Months Among All Participants: RFS was defined as the time between the date of randomization and the date of first melanoma recurrence (local, regional, distant metastasis) or death (whatever the cause), whichever occurred first. For participants who remained alive and whose disease had not recurred, RFS was censored on the date of last visit/contact with disease assessments. The percentage of participants with RFS at Month 6 was reported for all participants in both treatment arms of Part 1. 2. Part 1: Percentage of Participants With Recurrence-Free Survival (RFS) At 6 Months Among Participants With PD-L1-positive Tumor Expression: RFS was defined as the time between the date of randomization and the date of first melanoma recurrence (local, regional, distant metastasis) or death (whatever the cause), whichever occurred first. For participants who remained alive and whose disease had not recurred, RFS was censored on the date of last visit/contact with disease assessments. The percentage of participants with RFS at Month 6 was reported for all participants with PD-L1-positive tumors in both treatment arms of Part 1.
Secondary Obj.:	1. Distant Metastases-free Survival (DMFS) in All Participants 2. Distant Metastases-free Survival (DMFS) for Participants With PD-L1-positive Tumor Expression 3. Overall Survival (OS) for All Participants

2.2 Background & Rationale

This study will assess whether post-surgery therapy with pembrolizumab improves recurrence-free survival (RFS) as compared to placebo for high-risk participants with melanoma (Stage IIIA [≥ 1 mm metastasis], IIIB and IIIC). The study will also assess whether pembrolizumab improves RFS versus placebo in the subgroup of participants with

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

programmed cell death-ligand 1 (PD-L1)-positive tumor expression. Participants will be stratified for stage of disease and region and then will be randomly assigned to receive either pembrolizumab or placebo as post-surgery therapy in Part 1. In Part 2, participants who experience a disease recurrence are eligible for pembrolizumab treatment (if treated with placebo in Part 1) or pembrolizumab rechallenge (if treated with pembrolizumab in Part 1).

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE3, Status: ACTIVE_NOT_RECRUITING
Target N: 1019

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

Not Provided

4.2 Exclusion Criteria

Not Provided

11. Study Identifiers & Governance

Primary ID: 1137004121105929006
Registry Number: NCT02362594
Sponsor: Merck
Study Title: Study of Pembrolizumab (MK-3475) Versus Placebo After Complete Resection of High-Risk Stage III Melanoma (MK-3475-054/1325-MG/KEYNOTE-054)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____