

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase III, Randomized, Double-Blind, Placebo-Controlled Study Evaluating the Efficacy and Safety of Inavolisib Plus Palbociclib and Fulvestrant Versus Placebo Plus Palbociclib and Fulvestrant in Patients With PIK3CA-Mutant, Hormone Receptor-Positive, HER2-Negative, Locally Advanced or Metastatic Breast Cancer
Short Title:	A Study Evaluating the Efficacy and Safety of Inavolisib + Palbociclib + Fulvestrant vs Placebo + Palbociclib + Fulvestrant in Participants With PIK3CA-Mutant, Hormone Receptor-Positive, HER2-Negative, Locally Advanced or Metastatic Breast Cancer
Protocol Number:	188928349158633105
NCT Number:	NCT04191499
Sponsor Name:	Roche
Investigational Product:	fulvestrant
Phase:	PHASE2, PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Progression-Free Survival (PFS): PFS was defined as the time from randomization to the first occurrence of disease progression, as determined by the investigator according to Response Evaluation Criteria in Solid Tumors (RECIST), Version 1.1 or death from any cause (whichever occurs first). Progressive disease (PD) was defined as at least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum in the study, including baseline, in addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 millimeter (mm). The appearance of one or more new lesions was also considered progression. Data for participants without the occurrence of PD or death as of the clinical cutoff date (CCOD) were censored at the time of the last tumor assessment prior to the CCOD. Median PFS was calculated using the Kaplan-Meier methodology.
Secondary Obj.:	1. Percentage of Participants With Objective Response Rate (ORR) 2. Percentage of Participants With Best Overall Response Rate (BOR) 3. Duration of Response (DOR)

2.2 Background & Rationale

This study will evaluate the efficacy, safety, and pharmacokinetics of inavolisib in combination with palbociclib and fulvestrant compared with placebo plus palbociclib and fulvestrant in participants with PIK3CA-mutant, hormone receptor (HR)-positive, HER2-negative locally advanced or metastatic breast cancer whose disease progressed during treatment or within 12 months of completing adjuvant endocrine therapy and who have not received prior systemic therapy for metastatic disease.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE2, PHASE3, Status: ACTIVE_NOT_RECRUITING

Target N: 325

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

Not Provided

4.2 Exclusion Criteria

Not Provided

11. Study Identifiers & Governance

Primary ID: 188928349158633105

Registry Number: NCT04191499

Sponsor: Roche

Study Title: A Study Evaluating the Efficacy and Safety of Inavolisib + Palbociclib + Fulvestrant vs Placebo + Palbociclib + Fulvestrant in Participants With PIK3CA-Mutant, Hormone Receptor-Positive, HER2-Negative, Locally Advanced or Metastatic Breast Cancer

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____