

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 3, Randomized, Active-controlled, Open-label, Multicenter Study to Compare the Efficacy and Safety of MK-2870 Monotherapy Versus Treatment of Physician's Choice in Participants With Endometrial Cancer Who Have Received Prior Platinum-based Chemotherapy and Immunotherapy (MK-2870-005/ENGOT-en23/GOG-3095)
Short Title:	Sacituzumab Tirumotecan (MK-2870) in Post Platinum and Post Immunotherapy Endometrial Cancer (MK-2870-005)
Protocol Number:	36583125590353280
NCT Number:	NCT06132958
Sponsor Name:	Merck
Investigational Product:	doxorubicin
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Progression-free Survival (PFS) per Response Evaluation Criteria in Solid Tumors (RECIST) 1.1 as Assessed by Blinded Independent Central Review (BICR): PFS is defined as the time from randomization to the first documented disease progression per RECIST 1.1 based on BICR or death due to any cause, whichever occurs first. 2. Overall Survival (OS): OS is defined as the time from randomization to death due to any cause.
Secondary Obj.:	1. Objective Response Rate (ORR) per RECIST 1.1 as Assessed by BICR 2. Duration of Response (DOR) per RECIST 1.1 as Assessed by BICR 3. Number of Participants Who Experience One or More Adverse Events (AEs)

2.2 Background & Rationale

Researchers are looking for new ways to treat people with endometrial cancer (EC) who have previously received treatment with platinum based therapy (a type of chemotherapy) and immunotherapy. Immunotherapy is a treatment that helps the immune system fight cancer. This clinical study will compare sacituzumab tirumotecan to chemotherapy. The goal of the study is to learn if people who receive sacituzumab tirumotecan live longer overall and without the cancer getting worse compared to people who receive chemotherapy.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: ACTIVE_NOT_RECRUITING
Target N:	710

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Has a histologically-confirmed diagnosis of endometrial carcinoma or carcinosarcoma.
- * Has radiographically evaluable disease, either measurable or nonmeasurable per response evaluation criteria in solid tumors (RECIST 1.1), as assessed by blinded independent central review (BICR).
- * Has received prior platinum-based chemotherapy and anti-programmed cell death 1 protein (PD-1)/anti- programmed cell death ligand 1 (PD-L1) therapy, either separately or in combination.

4.2 Exclusion Criteria

- * Has neuroendocrine tumors or endometrial sarcoma, including stromal sarcoma, leiomyosarcoma, adenosarcoma, or other types of pure sarcomas
- * Has a history of documented severe dry eye syndrome, severe Meibomian gland disease and/or blepharitis, or severe corneal disease that prevents/delays corneal healing
- * Has active inflammatory bowel disease requiring immunosuppressive medication or previous history of inflammatory bowel disease
- * Has had a recurrence of endometrial carcinoma or carcinosarcoma more than >12 months after completing platinum-based therapy administered in the curative-intent setting without any additional platinum-based therapy received in the recurrent setting. Note: 1) If Immunotherapy-based treatment is administered in the recurrent setting, then platinum rechallenge is not required, regardless of the duration of the platinum-free interval from time of adjuvant therapy 2) For Stage IVb disease, treatment that includes gynecological surgery followed by a platinum-based regimen is NOT considered curative-intent per protocol and does not require platinum rechallenge in the recurrent setting, regardless of the duration of the platinum-free interval
- * Has received more than 3 prior lines of therapy for endometrial carcinoma or carcinosarcoma
- * Has history of (noninfectious) pneumonitis/interstitial lung disease that required steroids or has current pneumonitis/interstitial lung disease
- * Has received prior treatment with single-agent nonplatinum based chemotherapy in the third-line setting
- * Has received prior treatment with a trophoblast cell surface antigen 2 (TROP2)-targeted antibody drug conjugate (ADC) (eg, sacituzumab govitecan)
- * Has received prior treatment with a topoisomerase I inhibitor-containing ADC (eg, sacituzumab govitecan or fam-trastuzumab deruxtecan-nxki)
- * Has previously received both single-agent paclitaxel and single-agent doxorubicin in any setting for prior treatment of endometrial cancer
- * Requires recurrent drainage of effusions (e.g., pleural, ascitic, etc.) within 6 weeks before randomization

11. Study Identifiers & Governance

Primary ID:	36583125590353280
Registry Number:	NCT06132958
Sponsor:	Merck
Study Title:	Sacituzumab Tirumotecan (MK-2870) in Post Platinum and Post Immunotherapy Endometrial Cancer (MK-2870-005)

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____