

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	Phase 3, Open-label Trial to Evaluate Safety, Pharmacokinetics, and Efficacy of Benralizumab in Children With Eosinophilic Diseases (CLIPS)
Short Title:	A Trial to Investigate Benralizumab in Children With Eosinophilic Diseases
Protocol Number:	478362929341363011
NCT Number:	NCT06512883
Sponsor Name:	AstraZeneca
Investigational Product:	benralizumab
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Number of Participants with Adverse Events (AEs): The safety and tolerability of benralizumab will be evaluated. 2. Serum Concentrations of Benralizumab: The PK of benralizumab will be evaluated.
Secondary Obj.:	1. EGPA Cohort: Percentage of Participants with Remission at Week 24 2. Number of Participants with Positive Antidrug Antibody (ADA) 3. Change From Baseline in Peripheral Blood Eosinophil Count

2.2 Background & Rationale

The main purpose of study is to assess the safety, tolerability, and pharmacokinetic (PK) of benralizumab.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: RECRUITING
Target N:	14

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

All Cohorts:

- * Male or female patients must be aged 6 to < 18 years of age at the time of signing the assent form and their caregiver signing the informed consent form.
- * Body weight greater than ($\geq$) 15 kilograms (kg).

EGPA Cohort:

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

* Therapy with corticosteroids: The prescribed dose of oral corticosteroids (OCS) (greater than 0.1 milligrams per kilogram per day (mg/kg/day), max dose of 50 milligrams per day (mg/day) must be stable (that is, no adjustment of the dose) for at least 4 weeks prior to baseline (Visit 2).

* Immunosuppressive therapy: If receiving immunosuppressive therapy, the dosage must be stable for at least 4 weeks prior to baseline (Visit 2).

4.2 Exclusion Criteria

All Cohorts:

* Any current malignancy or history of malignancy.

* History of anaphylaxis to any biologic therapy or vaccine.

* Known, pre-existing, clinically significant endocrine, autoimmune, metabolic, neurological, renal, gastrointestinal, hepatic, haematological, respiratory, or any other system abnormalities.

* Previous receipt of benralizumab in an interventional clinical study.

EGPA Cohort:

* Diagnosed with granulomatosis with polyangiitis (previously known as Wegener's granulomatosis) or microscopic polyangiitis.

* EGPA relapse: any deterioration in EGPA and/or organ-threatening EGPA that per Investigator judgement renders patients unstable in their EGPA within 3 months prior to screening (Visit 1) and through first administration of IP at baseline (Visit 2).

* Life-threatening EGPA: imminently life-threatening EGPA disease within 3 months prior to screening (Visit 1) and through first administration of IP at baseline (Visit 2), as per Investigator judgement.

11. Study Identifiers & Governance

Primary ID: 478362929341363011

Registry Number: NCT06512883

Sponsor: AstraZeneca

Study Title: A Trial to Investigate Benralizumab in Children With Eosinophilic Diseases

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____