

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase III, Open-label, Randomised Study of Osimertinib With or Without Datopotamab Deruxtecan (Dato-DXd), as First-line Treatment in Participants With Epidermal Growth Factor Receptor (EGFR) Mutation-positive, Locally Advanced or Metastatic Non-small Cell Lung Cancer
Short Title:	Phase III, Open-label Study of First-line Osimertinib With or Without Datopotamab Deruxtecan for EGFRm Locally Advanced or Metastatic Non-small Cell Lung Cancer
Protocol Number:	51139092551510042
NCT Number:	NCT06350097
Sponsor Name:	AstraZeneca
Investigational Product:	osimertinib
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. To demonstrate the superiority of osimertinib in combination with Datopotamab Deruxtecan relative to osimertinib by assessment of Progression Free Survival (PFS) by BICR in all randomised participants.: PFS is defined as time from randomisation until progression per RECIST 1.1 as assessed by BICR, or death due to any cause (in the absence of progression).
Secondary Obj.:	1. To demonstrate the superiority of osimertinib in combination with Datopotamab Deruxtecan relative to osimertinib by assessment of Overall Survival (OS) in all randomised participants. 2. To demonstrate the effectiveness of osimertinib in combination with Datopotamab Deruxtecan relative to osimertinib by assessment of PFS on Central nervous system (CNS) metastases in participants with CNS metastases at baseline 3. To demonstrate the effectiveness of osimertinib in combination with Datopotamab Deruxtecan relative to osimertinib by assessment of PFS by investigator in all randomised participants.

2.2 Background & Rationale

The purpose of this study is to evaluate efficacy and safety of osimertinib (tablet) in combination with Dato-DXd (i.v. infusion) compared with osimertinib (tablet) monotherapy as a first-line therapy in participants with locally advanced or metastatic EGFRm (Ex19del and/or L858R) NSCLC.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: RECRUITING
Target N:	582

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

Age

1. Participant must be $\geq$ 18 years.

Type of Participant and Disease Characteristics

2. Histologically or cytologically documented nonsquamous NSCLC. NSCLC of mixed histology is allowed if adenocarcinoma is the predominant histology. Mixed small-cell lung cancer and NSCLC histology, and sarcomatoid variant of NSCLC is ineligible.
3. Stage IIIB or IIIC or Stage IV metastatic NSCLC or recurrent NSCLC (based on the American Joint Committee on Cancer Edition 8) not amenable to curative surgery or definitive chemoradiation at the time of randomisation.
4. Participants must not have received prior EGFR TKIs or other systemic therapy for Stage IIIB, IIIC or IV NSCLC.
5. The tumour harbors at least 1 of the 2 common EGFR mutations known to be associated with EGFR-TKI sensitivity (Ex19del or L858R), either alone or in combination with other genomic alterations, which may include EGFR T790M, assessed by a CLIA-certified (US sites) or an accredited (outside of the US) local laboratory or by central prospective tissue testing.
6. For participants enrolled in randomisation period, mandatory provision of an unstained, archival tumour tissue sample in a quantity sufficient to allow for central confirmation of the EGFR mutation status.
7. WHO performance status of 0 or 1.
8. At least one lesion not previously irradiated that qualifies as a RECIST 1.1 TL at baseline and can be accurately measured at baseline as $\geq$ 10 mm in the longest diameter (except lymph nodes, which must have short axis $\geq$ 15 mm) with CT or MRI and is suitable for accurate repeated measurements.
9. Adequate bone marrow reserve and organ function within 7 days before the first dose of study intervention.

Sex and Contraceptive/Barrier Requirements

10. Contraceptive use by males or females should be consistent with local regulations regarding the methods of contraception for those participating in clinical studies.

Other Inclusion Criteria

11. All races, gender and ethnic groups are eligible for this study.

4.2 Exclusion Criteria

1. As judged by the investigator, any evidence of diseases (such as severe or uncontrolled systemic diseases, including active bleeding diseases, psychiatric illness/social situations), history of allogenic organ transplant, and/or substance abuse which, in the investigator's opinion, makes it undesirable for the participant to participate in the study or that would jeopardise compliance with the protocol.
2. Refractory nausea and vomiting, chronic gastrointestinal disease, inability to swallow the formulated product, or previous significant bowel resection that would preclude adequate absorption, distribution, metabolism, or excretion of osimertinib.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

3. History of another primary malignancy.
4. Spinal cord compression and unstable brain metastases, as defined by Protocol.
5. Clinically significant corneal disease.
6. Has active or uncontrolled hepatitis B or C virus infection, as defined by Protocol.
7. Known HIV infection that is not well controlled, as defined by Protocol.
8. Uncontrolled infection requiring i.v. antibiotics, antivirals or antifungals; suspected infections (eg, prodromal symptoms); or inability to rule out infections (participants with localised fungal infections of skin or nails are eligible).
9. Resting ECG with clinically abnormal findings, as defined by Protocol.
10. Uncontrolled or significant cardiac disease, as defined by Protocol.
11. Past medical history of ILD/pneumonitis, including radiation pneumonitis (apart from radiation pneumonitis that did not require steroids), or drug-induced ILD/pneumonitis, or has suspected ILD/pneumonitis that cannot be ruled out by imaging at screening.
12. Pulmonary embolism within 3 months of the study enrolment or has severe pulmonary function compromise.

Prior/Concomitant Therapy

13. Prior exposure to any agent including an ADC containing a chemotherapeutic agent targeting topoisomerase I, TROP2-targeted therapy.

Prior/Concurrent Clinical Study Experience

14. Participants with a known history of severe hypersensitivity reactions to either Dato-DXd and osimertinib or any excipients of Dato DXd and osimertinib or drugs with a similar chemical structure or class to DXd and osimertinib.

11. Study Identifiers & Governance

Primary ID:	51139092551510042
Registry Number:	NCT06350097
Sponsor:	AstraZeneca
Study Title:	Phase III, Open-label Study of First-line Osimertinib With or Without Datopotamab Deruxtecan for EGFRm Locally Advanced or Metastatic Non-small Cell Lung Cancer

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____