

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase Ib/III Randomized, Multicenter, Global Study of Volrustomig Plus Casdatifan or Volrustomig Monotherapy Versus Nivolumab Plus Ipilimumab as First-line Treatment for Participants With Advanced Clear Cell Renal Cell Carcinoma (ccRCC)
Short Title:	A Study to Investigate the Efficacy and Safety of Volrustomig ± Casdatifan vs Nivolumab + Ipilimumab as 1L Treatment for Advanced ccRCC
Protocol Number:	764043096871782188
NCT Number:	NCT07000149
Sponsor Name:	AstraZeneca
Investigational Product:	ipilimumab
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Phase Ib: Number of participants with adverse events (AEs) and serious adverse events (SAEs): Number of participants who received at least one dose of study treatment will be assessed. 2. Phase III: Progression-free Survival (PFS): The PFS is defined as the time from randomization until radiological progression or death due to any cause (in the absence of progression). 3. Phase III: Overall Survival (OS): The OS is defined as the time from randomization until the date of death due to any cause.
Secondary Obj.:	1. Phase Ib: Objective Response rate (ORR) 2. Phase Ib: Duration of Response (DoR) 3. Phase Ib: Progression-free Survival (PFS)

2.2 Background & Rationale

This is a Phase Ib/III, randomized, multicenter, global study evaluating the efficacy and safety of volrustomig in combination with casdatifan for the first-line (1L) treatment of participants with advanced clear cell renal cell carcinoma (ccRCC).

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: ACTIVE_NOT_RECRUITING
Target N:	1116

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Histologically or cytologically confirmed RCC with clear cell component.
- * Advanced/metastatic RCC or recurrent disease that has not previously been treated with systemic therapy in the 1L

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

setting.

- * Karnofsky Performance Status ? 70%.

- * Provision of acceptable tumor sample.

- * At least one lesion that can be accurately measured at baseline as ? 10 mm in the longest diameter (except lymph nodes, which must have short axis ? 15 mm) with computed tomography (CT) or magnetic resonance imaging (MRI) and is suitable for accurate repeated measurements.

4.2 Exclusion Criteria

- * History of leptomeningeal disease or spinal cord compression.

- * Symptomatic brain metastases.

- * Medical history of severe chronic obstructive pulmonary disease.

- * Active or prior documented autoimmune or inflammatory disorders.

- * Prior systemic therapy for advanced/metastatic RCC. Note - Other inclusion and exclusion criteria may apply as stated in the protocol.

11. Study Identifiers & Governance

Primary ID: 764043096871782188

Registry Number: NCT07000149

Sponsor: AstraZeneca

Study Title: A Study to Investigate the Efficacy and Safety of Volrustomig ± Casdatifan vs Nivolumab + Ipilimumab as 1L Treatment for Advanced ccRCC

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____