

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 3 Study of MK-4280A (Coformulated Favezelimab [MK-4280] Plus Pembrolizumab [MK-3475]) Versus Standard of Care in Previously Treated Metastatic PD-L1 Positive Colorectal Cancer (KEYFORM-007)
Short Title:	A Study of Coformulated Favezelimab/Pembrolizumab (MK-4280A) Versus Standard of Care in Subjects With Previously Treated Metastatic PD-L1 Positive Colorectal Cancer (MK-4280A-007)-China Extension Study
Protocol Number:	865249763829750483
NCT Number:	NCT05600309
Sponsor Name:	Merck
Investigational Product:	regorafenib
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Overall Survival (OS): OS was defined as the time from randomization to death due to any cause.
Secondary Obj.:	1. Progression-Free Survival (PFS) According Per Response Evaluation Criteria in Solid Tumors Version 1.1 (RECIST 1.1) as Assessed by Blinded Independent Central Review (BICR) 2. Objective Response Rate (ORR) Per RECIST 1.1 as Assessed by BICR 3. Duration of Response (DOR) Per RECIST 1.1 as Assessed by BICR

2.2 Background & Rationale

The purpose of this China extension study is to assess the safety and efficacy of coformulated favezelimab/pembrolizumab (MK-4280A) in adult Chinese participants with metastatic colorectal cancer. The study will also compare MK-4280A with the standard of care treatment of regorafenib and TAS-102 (trifluridine and tipiracil).

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: COMPLETED
Target N:	94

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Has a histologically confirmed colorectal adenocarcinoma that is metastatic and unresectable.
- * Has measurable disease per RECIST 1.1 as assessed by the local site investigator.
- * Has been previously treated for the disease and radiographically progressed on or after or could not tolerate standard

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

treatment.

- * Submits an archival (? 5 years) or newly obtained tumor tissue sample or newly obtained tumor tissue sample that has not been previously irradiated.
- * Has an Eastern Cooperative Oncology Group Performance Score (ECOG PS) of 0 to 1 within 10 days prior to first dose of study intervention.
- * Has a life expectancy of at least 3 months, based on the investigator assessment.
- * Has the ability to swallow and retain oral medication and not have any clinically significant gastrointestinal abnormalities that might alter absorption.
- * Has adequate organ function.

4.2 Exclusion Criteria

- * Has previously been found to have deficient mismatch repair/microsatellite instability-high (dMMR/MSI-H) tumor status.
- * Has known active central nervous system (CNS) metastases and/or carcinomatous meningitis or leptomeningeal disease.
- * Has a history of acute or chronic pancreatitis.
- * Has neuromuscular disorders associated with an elevated creatine kinase (eg, inflammatory myopathies, muscular dystrophy, amyotrophic lateral sclerosis, spinal muscular atrophy)
- * Has clinically significant cardiovascular disease within 12 months from first dose of study intervention, including New York Heart Association Class III or IV congestive heart failure, unstable angina, myocardial infarction, cerebral vascular accident, or cardiac arrhythmia associated with hemodynamic instability.
- * Has urine protein greater than or equal to 1g/24h.
- * A woman of childbearing potential who has a positive urine/serum pregnancy test within 24/72 hours prior to the first dose of study intervention.
- * Has received prior therapy with an anti-programmed cell death 1 (PD-1), anti-programmed death ligand 1 (PD-L1), or anti-programmed cell death ligand 2 (PD-L2), anti-lymphocyte activation gene 3 (LAG-3) antibody, with a tyrosine kinase inhibitor (TKI; eg, lenvatinib) other than rapidly accelerated fibrosarcoma (RAF) inhibitors (binimetinib is permitted if combined with a RAF inhibitor), or with an agent directed to another stimulatory or coinhibitory T-cell receptor (eg, cytotoxic T-lymphocyte-associated protein 4, OX-40, cluster of differentiation [CD] 137).
- * Has previously received regorafenib or TAS-102.
- * Has received prior systemic anticancer therapy including investigational agents within 28 days before randomization.
- * Has received prior radiotherapy within 2 weeks of start of study intervention. Participants must have recovered from all radiation-related toxicities, not require corticosteroids, and not have had radiation pneumonitis. A 1-week washout is permitted for palliative radiation (?2 weeks of radiotherapy) to non-CNS disease.
- * Has received a live or live-attenuated vaccine within 30 days before the first dose of study intervention.
- * Has a known additional malignancy that is progressing or has required active treatment within the past 3 years.
- * Has severe hypersensitivity (?Grade 3) to pembrolizumab and/or any of its excipients.
- * Has an active autoimmune disease that has required systemic treatment in past 2 years.
- * Has a history of (noninfectious) pneumonitis/interstitial lung disease that required steroids or has current pneumonitis/interstitial lung disease.
- * Has an active infection requiring systemic therapy (eg, tuberculosis, known viral or bacterial infections, etc.).
- * Has a known history of human immunodeficiency virus (HIV) infection.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Has known history of Hepatitis B or known active Hepatitis C virus infection.
- * Has a history or current evidence of any condition, therapy, or laboratory abnormality that might confound the results of the study, interfere with the participant's participation for the full duration of the study, or is not in the best interest of the participant to participate, in the opinion of the treating investigator.
- * Has had an allogenic tissue/solid organ transplant.

11. Study Identifiers & Governance

Primary ID: 865249763829750483

Registry Number: NCT05600309

Sponsor: Merck

Study Title: A Study of Coformulated Favezelimab/Pembrolizumab (MK-4280A) Versus Standard of Care in Subjects With Previously Treated Metastatic PD-L1 Positive Colorectal Cancer (MK-4280A-007)-China Extension Study

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____