

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 2, Randomized, Double-Blind, Placebo-controlled Study to Compare Efficacy and Safety of Oral Azacitidine Plus Best Supportive Care Versus Best Supportive Care as Maintenance Therapy in Japanese Subjects With Acute Myeloid Leukemia in Complete Remission
Short Title:	A Study to Compare the Efficacy and Safety of Oral Azacitidine Plus Best Supportive Care Versus Best Supportive Care as Maintenance Therapy in Japanese Participants With Acute Myeloid Leukemia (AML) in Complete Remission
Protocol Number:	1038523475209203280
NCT Number:	NCT05197426
Sponsor Name:	Bristol Myers Squibb
Investigational Product:	azacitidine
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Relapse-free Survival (RFS)
Secondary Obj.:	1. Overall Survival (OS) 2. Time to relapse from Complete Remission (CR) 3. Time to relapse from complete remission with incomplete blood count recovery (CRi)

2.2 Background & Rationale

The purpose of this study is to assess the efficacy and safety of oral azacitidine plus best supportive care versus best supportive care as maintenance therapy in a cohort of Japanese participants ≥ 55 years of age with Acute Myeloid Leukemia (AML) and in complete remission/complete remission with incomplete blood count recovery after conventional induction chemotherapy with or without consolidation chemotherapy.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE2, Status: ACTIVE_NOT_RECRUITING
Target N:	19

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * ≥ 55 years of age inclusive at the time of signing the informed consent
- * Newly diagnosed, histologically confirmed de novo Acute Myeloid Leukemia (AML) or AML secondary to prior myelodysplastic syndrome (MDS) or chronic myelomonocytic leukemia (CMML)
- * Should have undergone induction therapy with intensive chemotherapy with or without consolidation therapy as

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

recommended in appropriate guideline(s) or equivalent regimen according to institutional standard: having achieved first complete remission (CR)/complete remission with incomplete blood count recovery (CRi) status within 4 months prior to starting study therapy

4.2 Exclusion Criteria

- * Suspected or proven acute promyelocytic leukemia; or AML with previous hematologic disorder such as chronic myeloid leukemia or myeloproliferative neoplasms, excluding MDS and CMML
- * Prior bone marrow or stem cell transplantation
- * Received therapy with hypomethylating agents for MDS and went on to develop AML within four months of discontinuing the therapy with hypomethylating agents
- * Have achieved CR/CRi following therapy with hypomethylating agents

Other protocol-defined inclusion/exclusion criteria apply

11. Study Identifiers & Governance

Primary ID:	1038523475209203280
Registry Number:	NCT05197426
Sponsor:	Bristol Myers Squibb
Study Title:	A Study to Compare the Efficacy and Safety of Oral Azacitidine Plus Best Supportive Care Versus Best Supportive Care as Maintenance Therapy in Japanese Participants With Acute Myeloid Leukemia (AML) in Complete Remission

15. Sign-off & Approval

Principal Investigator: _____	Date: _____
Clinical Operations Lead: _____	Date: _____
Lead Statistician: _____	Date: _____