

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A PHASE 1B/2, OPEN-LABEL STUDY TO EVALUATE THE SAFETY, PHARMACOKINETICS, PHARMACODYNAMICS, AND EFFICACY OF ELRANATAMAB (PF-06863135) IN CHINESE PARTICIPANTS WITH MULTIPLE MYELOMA WHO ARE REFRACTORY TO AT LEAST ONE PROTEASOME INHIBITOR, ONE IMMUNOMODULATORY DRUG AND ONE ANTI-CD38 ANTIBODY (TRIPLE-CLASS REFRACTORY MM)
Short Title:	A Study of Elranatamab (PF-06863135) in Chinese Participants With Refractory Multiple Myeloma.
Protocol Number:	1063046164827300209
NCT Number:	NCT05228470
Sponsor Name:	Pfizer
Investigational Product:	elranatamab
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Phase 1b: Number of Participants With Dose-Limiting Toxicities (DLT): Grade(G)4 neutropenia ≥ 5 day; febrile neutropenia (absolute neutrophil count $[ANC] < 1000/\text{millimeter cube (mm}^3)$ with single temperature ≥ 38.3 degree Celsius (deg C) or sustained temp ≥ 38 deg C for ≥ 1 hour(H); $G \geq 3$ neutropenia with infection; G4 thrombocytopenia (unless baseline count $\geq 25,000/\text{mm}^3$ and $< 50,000/\text{mm}^3$, in which case G4 thrombocytopenia to be accompanied by $\geq G2$ bleeding); Platelet count $< 10,000/\text{mm}^3$; G3 thrombocytopenia with $\geq G2$ bleeding; $G \geq 4$ AE; G3 cytokine release syndrome(CRS) except CRS not maximally treated/improved to $\leq G1$ within 48H; G3 AE except AE attributed to CRS, G3 nausea,vomiting,diarrhea improved to $G \leq 2$ within 72H after medical management, G3 fatigue < 1 week, G3 AE recovered to baseline/G1 within 5 day; confirmed drug-induced liver injury; G3-4 laboratory (lab) abnormality except G3-4 lab abnormality improved to $G \leq 2$ within 72H after medical management $\&$ without sequelae;G3 injection site reaction; G2/other clinically important AE may be considered DLT. 2. Objective Response Rate (ORR) by Blinded Independent Central Review (BICR) as Per International Myeloma Working Group (IMWG) Criteria: ORR: Percentage of participants with best overall response of confirmed stringent complete response (sCR), CR, very good partial response (VGPR) or PR per IMWG criteria. sCR: CR $\&$ normal serum free light chain (sFLC) ratio $\&$ absence of clonal cells in BMB/BMA by IH, IF, or flow cytometry. CR: negative immunofixation on serum $\&$ urine, disappearance of any soft tissue plasmacytoma $\&$
--------------------	--

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

<5% plasma cells in BMA, if disease measurable by sFLC only, preceding criteria plus normal sFLC ratio. VGPR: Serum & urine M-protein detectable by immunofixation but not on electrophoresis; or >=90% reduction in serum M-protein & urine M-protein level <100mg/24h. PR: >=50% reduction in serum M-protein & reduction in 24h urinary M-protein by >=90% or <200 mg/24h. If serum & urine M-protein were unmeasurable, VGPR & PR: >=90% & >=50% decrease in difference respectively between involved & uninvolved sFLC levels & if present at baseline, >=90% & >=50% reduction in soft tissue plasmacytomas' size.

Secondary Obj.:

1. Duration of Response (DOR) as Per IMWG Criteria by BICR 2. Duration of Response (DOR) as Per IMWG Criteria by Investigator Assessment 3. Complete Response Rate (CRR) as Per IMWG Criteria by BICR

2.2 Background & Rationale

The purpose of this study is to understand the study medicine (called Elranatamab, or PF-06863135) as potential treatment for refractory multiple myeloma. Multiple myeloma is a form of cancer in the bone that forces healthy blood cells to go out. Sometimes, multiple myeloma does not respond to current therapy or quickly progresses, and this is called refractory multiple myeloma.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE2, Status: ACTIVE_NOT_RECRUITING

Target N: 39

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Diagnosis of multiple myeloma (IMWG criteria, Rajkumar et al, 2014)
- * Measurable disease, as defined by at least 1 of the following:
 - * Serum M-protein ≥0.5 g/dL
 - * Urinary M-protein excretion ≥200 mg/24 hours
 - * Serum immunoglobulin FLC ≥10 mg/dL (≥100 mg/L) AND abnormal serum immunoglobulin kappa to lambda FLC ratio
- * Refractory to at least one IMiD
- * Refractory to at least one PI
- * Refractory to at least one anti-CD38 antibody
- * Relapsed/refractory to last anti-myeloma regimen
- * ECOG performance status ≤2
- * Adequate BM function characterized by the following:
 1. Absolute neutrophil count ≥1.0 × 10⁹/L
 2. Platelets ≥ 25 × 10⁹/L
 3. Hemoglobin ≥8 g/dL
- * Resolved acute effects of any prior therapy to baseline severity or CTCAE Grade ≤1

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

* Not pregnant and willing to use contraception

4.2 Exclusion Criteria

- * Smoldering multiple myeloma
- * Active Plasma cell leukemia
- * Amyloidosis
- * POEMS syndrome
- * Stem cell transplant or active GVHD within 12 weeks prior to enrollment.
- * Previous treatment with an anti-BCMA directed therapy
- * Impaired cardiovascular function or clinically significant cardiovascular diseases
- * Ongoing Grade ?2 peripheral sensory or motor neuropathy. History of GBS or GBS variants, or history of any Grade ?3 peripheral motor polyneuropathy.
- * Active HBV, HCV, SARS-CoV2, HIV, or any active, uncontrolled bacterial, fungal, or viral infection
- * Any other active malignancy within 3 years prior to enrollment, except for adequately treated basal cell or squamous cell skin cancer, or carcinoma in situ.
- * Previous administration with an investigational drug within 30 days or 5 half-lives preceding the first dose of study intervention used in this study (whichever is longer)

11. Study Identifiers & Governance

Primary ID: 1063046164827300209

Registry Number: NCT05228470

Sponsor: Pfizer

Study Title: A Study of Elranatamab (PF-06863135) in Chinese Participants With Refractory Multiple Myeloma.

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____