

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 2, Open-label, Study to Evaluate the Efficacy and Safety of Rapcabtagene Autoleucel in Patients With Active, Refractory Systemic Lupus Erythematosus (SLE) or Active, Refractory Lupus Nephritis (LN).
Short Title:	A Study of Rapcabtagene Autoleucel in Active, Refractory Systemic Lupus Erythematosus (SLE) or Lupus Nephritis (LN) Patients (AUTOGRAPH - SLE/LN)
Protocol Number:	179478236634708606
NCT Number:	NCT06581198
Sponsor Name:	Novartis
Investigational Product:	ciltacabtagene autoleucel
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Evaluate the efficacy of rapcabtagene autoleucel: Defined as: Meeting the criteria of the Definition Of Remission In Systemic Lupus Erythematosus (DORIS) or Achieving complete renal response (CRR)
Secondary Obj.:	1. Percentage of participants achieving complete renal response (CRR) 2. Number of weeks where Lupus Low Disease Activity Score (LLDAS) was achieved 3. PPercentage of participants without flaring (i.e., 1 new BILAG2004 A or 2 new BILAG2004 B flares)

2.2 Background & Rationale

The purpose of this study is to evaluate the efficacy and safety of rapcabtagene autoleucel (administered once following lymphodepletion) in patients with active, refractory systemic lupus erythematosus (SLE) or active, refractory lupus nephritis (LN).

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE2, Status: RECRUITING
Target N:	179

4. Subject Selection (Eligibility Criteria)

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4.1 Inclusion Criteria

- * Men and women with SLE, aged ≥ 18 years and ≤ 75 years at screening, fulfilling the 2019 European League Against Rheumatism (EULAR)/American College of Rheumatology (ACR) classification criteria for SLE at screening.
- * Participant must be positive for at least one of the following autoantibodies at screening: antinuclear antibodies (ANA) at a titer of $\geq 1:80$ (on HEp-2 cells or an equivalent positive test), or anti-dsDNA (above the ULN); or anti-Sm (above the ULN) as determined by a central laboratory.
- * Active lupus nephritis without signs of significant chronicity or active systemic lupus erythematosus
- * SLEDAI-2K Criteria at screening: SLEDAI-2K score ≥ 6 points (Gladman et al 2002, Touma et al 2011), excluding points attributed to "fever", "lupus headache", "alopecia", and "organic brain syndrome".
- * Inadequate response at screening to at least two therapies

Key Exclusion Criteria:

- * Any acute, severe lupus related-flare at screening that needs immediate treatment other than pulse GCs and/or makes the immunosuppressive washout impossible and, thus, makes the participant ineligible for CD19 CAR-T therapy
- * Inadequate organ function during screening and prior to randomization
- * History or current diagnosis of ECG or cardiac abnormalities indicating significant risk of safety for participants prior to randomization
- * Human immunodeficiency virus (HIV) positivity at screening.
- * Acute or chronic infection with hepatitis B (HBV) or hepatitis C (HCV) at screening.
- * Grade 2 or higher thromboembolic event in the past 4 weeks prior to screening.

Other protocol-defined inclusion/exclusion criteria may apply.

4.2 Exclusion Criteria

- * Any acute, severe lupus related-flare at screening that needs immediate treatment other than pulse GCs and/or makes the immunosuppressive washout impossible and, thus, makes the participant ineligible for CD19 CAR-T therapy
- * Inadequate organ function during screening and prior to randomization
- * History or current diagnosis of ECG or cardiac abnormalities indicating significant risk of safety for participants prior to randomization
- * Human immunodeficiency virus (HIV) positivity at screening.
- * Acute or chronic infection with hepatitis B (HBV) or hepatitis C (HCV) at screening.
- * Grade 2 or higher thromboembolic event in the past 4 weeks prior to screening.

Other protocol-defined inclusion/exclusion criteria may apply.

11. Study Identifiers & Governance

Primary ID:	179478236634708606
Registry Number:	NCT06581198
Sponsor:	Novartis
Study Title:	A Study of Rapcabtagene Autoleucel in Active, Refractory Systemic Lupus

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Erythematosus (SLE) or Lupus Nephritis (LN) Patients (AUTOGRAPH - SLE/LN)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____