

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	An Open-Label, Parallel-Group Study to Evaluate Safety, Tolerability, Pharmacokinetics, and Pharmacodynamic Effects of Ocrelizumab in Children and Adolescents With Relapsing-Remitting Multiple Sclerosis
Short Title:	A Study of Ocrelizumab in Children and Adolescents With Relapsing-Remitting Multiple Sclerosis
Protocol Number:	207230974578718374
NCT Number:	NCT04075266
Sponsor Name:	Roche
Investigational Product:	ocrelizumab
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Dose Exploration Period: Area Under the Concentration Versus Time Curve of Ocrelizumab: The population PK model was used to simulate concentration-time course and predict individual area under the concentration versus time curve. The data for the PK parameter: area under the concentration versus time curve was collected and analyzed as per body weight range (≤ 40 kg to ≥ 40 kg). 2. Dose Exploration Period: Maximum Concentration (Cmax) of Ocrelizumab: The population PK model was used to simulate concentration-time course and predict individual Cmax. The data for the PK parameter: Cmax was collected and analyzed as per body weight range (≤ 40 kg to ≥ 40 kg). 3. Dose Exploration Period: Levels of CD 19+ B-cell Count in Blood
Secondary Obj.:	1. Number of Participants With Adverse Events (AEs) 2. Dose Exploration Period: Level of Circulating T Cells and Natural Killer (NK) Cells 3. OOE Period: Level of Circulating T Cells and NK Cells

2.2 Background & Rationale

This 2-year study will evaluate the safety, tolerability, pharmacokinetics, and pharmacodynamic (PD) effects of ocrelizumab in children and adolescents ages ≥ 10 to ≤ 18 years with relapsing-remitting multiple sclerosis (RRMS). The data from this study will serve to determine the dosing regimen of ocrelizumab to be further investigated in the subsequent Phase III study in children and adolescents.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE2, Status: ACTIVE_NOT_RECRUITING
Target N:	23

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Body weight ≥ 25 kg
- * Children and adolescents must have received all childhood required vaccinations
- * Female participants of childbearing potential must agree to either remain completely abstinent or to use reliable means of contraception
- * Diagnosis of relapsing-remitting multiple sclerosis (RRMS)
- * Expanded Disability Status Scale (EDSS) at screening: 0-5.5, inclusive
- * Neurologic stability for ≥ 30 days prior to screening, and between screening and baseline
- * Participants naive to prior disease-modifying therapy (DMT)
- * Participants who have had at least 6 contiguous months of DMT within the past 1 year must have evidence of disease activity occurring after the full 6-month course of treatment, that is, at least one relapse or ≥ 1 Gd-enhancing lesion(s) on a T1-weighted brain MRI

4.2 Exclusion Criteria

- * Known presence or suspicion of other neurologic disorders that may mimic MS, including, but not limited to, acute disseminated encephalomyelitis, neuromyelitis optica or neuromyelitis optica spectrum disorders and any neurologic, somatic, or metabolic condition that could interfere with brain function or normal cognitive or neurological development
- * Patients that are aquaporin 4 positive and myelin oligodendrocyte glycoprotein (MOG) antibody positive are not eligible to participate in the study.
- * In case of an ADEM-like appearance of the first MS attack, a second attack with clear MS-like features is required.
- * Infection requiring hospitalization or treatment with IV anti-infective agents
- * History or known presence of recurrent or chronic infection (e.g., HIV, syphilis, tuberculosis)
- * Receipt of a live or live-attenuated vaccine within 6 weeks prior to treatment allocation
- * History or laboratory evidence of coagulation disorders
- * Peripheral venous access that precludes IV administration and venous blood sampling
- * Inability to complete a magnetic resonance imaging (MRI) scan
- * History of cancer, including solid tumors, hematologic malignancies, and carcinoma in situ
- * History of a severe allergic or anaphylactic reaction to humanized or murine monoclonal antibody (mAbs) or known hypersensitivity to any component of ocrelizumab solution
- * Previous treatment with B-cell-targeted therapies
- * Percentage of CD4 $< 30\%$
- * Absolute Neutrophil Count $< 1.5 \times 1000/\text{microliter}$
- * Lymphocyte count below the lower limit of normal (LLN) for age- and sex-specific reference range

11. Study Identifiers & Governance

Primary ID:	207230974578718374
Registry Number:	NCT04075266
Sponsor:	Roche
Study Title:	A Study of Ocrelizumab in Children and Adolescents With Relapsing-Remitting Multiple

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Sclerosis

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____