

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 1b/2 Dose Escalation and Expansion Study of Tucatinib in Combination With Trastuzumab and Oxaliplatin-based Chemotherapy or Pembrolizumab-containing Combinations for HER2+ Gastrointestinal Cancers
Short Title:	Tucatinib Plus Trastuzumab and Oxaliplatin-based Chemotherapy or Pembrolizumab-containing Combinations for HER2+ Gastrointestinal Cancers
Protocol Number:	276836765530532585
NCT Number:	NCT04430738
Sponsor Name:	Pfizer
Investigational Product:	fluorouracil
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Number of Participants With Renal Dose-Limiting Toxicities (DLTs): Phase 1b (Cohort 1A and 1B): A renal DLT was defined as an increase in serum cystatin C $>1.5\times$ baseline that was not related to pre-renal or post-renal etiologies (including disease progression, dehydration and intercurrent illness) and occurred during the period of treatment with tucatinib in combination with trastuzumab and FOLFOX between the first dose of tucatinib and the end of Cycle 3. Increased serum cystatin C for which there was an alternative clinical explanation (eg, clearly related to an intercurrent illness or disease progression) was not considered renal DLTs. 2. Number of Participants With Treatment Emergent Adverse Events (TEAEs): Phase 1b (Cohort 1E and 1F): An adverse event (AE) was defined as any untoward medical occurrence in a clinical investigational participant administered a medicinal product and which did not necessarily have a causal relationship with this treatment. TEAE was defined as a newly occurring or worsening AE after the first dose of study treatment (tucatinib, trastuzumab, leucovorin, fluorouracil, oxaliplatin and pembrolizumab) through 30 days after the last dose of study treatment. 3. Number of Participants With Treatment Related TEAEs: Phase 1b (Cohort 1E and 1F): An AE was defined as any untoward medical occurrence in a clinical investigational participant administered a medicinal product and which did not necessarily have a causal relationship with this treatment. TEAE was defined as a newly occurring or worsening AE after the first dose of study treatment (tucatinib, trastuzumab, leucovorin, fluorouracil, oxaliplatin and pembrolizumab) through 30 days after the last dose of study treatment. Treatment related TEAEs were AEs related to any study treatment and relatedness was judged by investigator.
Secondary Obj.:	1. Number of Participants With TEAEs: Phase 1b (Cohort 1A and 1B) 2. Number of

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Participants With Treatment Related TEAEs: Phase 1b (Cohort 1A and 1B) 3. Number of Participants With $\geq$ Grade 3 TEAEs: Phase 1b (Cohort 1A and 1B)

2.2 Background & Rationale

This trial studies tucatinib to find out if it is safe when given with trastuzumab and other anti-cancer drugs (pembrolizumab, FOLFOX, and CAPOX). It will look at what side effects happen when participants take this combination of drugs. A side effect is anything the drug does other than treating cancer. It will also look at whether tucatinib works with these drugs to treat certain types of cancer.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE2, Status: ACTIVE_NOT_RECRUITING

Target N: 40

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

* Participants must have an unresectable or metastatic solid malignancy that is histologically or cytologically confirmed to be one of the tumor types listed below:

* Cohorts 1A, 1B, 1C, and 1D

* CRC

* Gastric adenocarcinoma

* GEJ adenocarcinoma

* Esophageal adenocarcinoma

* Cholangiocarcinoma

* Gallbladder carcinoma

* Cohorts 1E, 1F, 1G, and 2A

* Gastric adenocarcinoma

* GEJ adenocarcinoma

* Esophageal adenocarcinoma

* Cohort 2B

* CRC

* Participants must be candidates to receive an oxaliplatin-based regimen as part of their standard-of-care treatment for all cohorts, except Cohort 1G.

* HER2+ disease, as determined by historic or local laboratory testing

* Phase 1b cohorts: measurable or non-measurable disease according to RECIST v1.1 as determined by the investigator

* Phase 2 cohorts: measurable disease according to RECIST v1.1 as determined by the investigator

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

* Eastern Cooperative Oncology Group Performance Status score of 0 or 1.

4.2 Exclusion Criteria

* History of known hypersensitivity to planned study treatment

* Known to be positive for Hepatitis B or C

* For Cohorts 2A and 2B: prior anti-HER2 therapies

* For Cohorts 1E, 1F, 1G, 2A: Prior therapy with an anti-PD-1, anti-PD-L1, or anti PD L2 agent or with an agent directed to another stimulatory or co-inhibitory T-cell receptor (eg, CTLA-4, OX 40, CD137), and was discontinued from that treatment due to a Grade 3 or higher immune-related adverse event (irAE)

There are additional inclusion criteria. The study center will determine if criteria for participations are met.

11. Study Identifiers & Governance

Primary ID: 276836765530532585

Registry Number: NCT04430738

Sponsor: Pfizer

Study Title: Tucatinib Plus Trastuzumab and Oxaliplatin-based Chemotherapy or Pembrolizumab-containing Combinations for HER2+ Gastrointestinal Cancers

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____