

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase II, Single-arm Study of De-escalation and Treatment-free Remission in Patients With Chronic Myeloid Leukemia Treated With Nilotinib in First-line Therapy Followed by a Second Attempt After Nilotinib and Asciminib Combination: DANTE Study
Short Title:	A Study of Full Treatment-free Remission in Patients With Chronic Myeloid Leukemia
Protocol Number:	332181817887867375
NCT Number:	NCT03874858
Sponsor Name:	Novartis
Investigational Product:	nilotinib
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Percentage of patients in full treatment-free remission 96 weeks after the start of the consolidation period of TFR1 stage: Full Treatment-Free Remission (FTFR) is defined as patients with Major Molecular Response (MMR) or better (i.e. BCR-ABL $\leq$ 0.1% IS), including those who have totally discontinued treatment during the Treatment-Free Remission (TFR) stage and those who are treated with half the standard dosage. Percentage of patients in full treatment-free remission 96 weeks after the start of the consolidation period of TFR1 stage is calculated by dividing the number of patients with no loss of MMR-Major Molecular Response (BCR-ABL $\leq$ 0.1% (IS) after 96 weeks by the number of patients who entered consolidation of TFR1 stage. 2. Percentage of patients in treatment-free remission 48 weeks after starting a second attempt at treatment-free remission during TFR2 stage.: Treatment-Free Remission (TFR) is defined as patients with MMR or better (i.e. BCR-ABL $\leq$ 0.1% IS). Percentage of patients in TFR 48 weeks after the start of TFR2 phase during TFR2 stage is calculated by dividing the number of patients with no loss of MMR after 48 weeks by the number of patients who entered TFR2 during TFR2 stage. Patients who require re-initiation of nilotinib during TFR2 for loss of MMR, and those discontinued from the study will be considered failures.
Secondary Obj.:	1. Percentage of patients who remain in sustained Deep Molecular Response (DMR) at the end of the consolidation period (week 48 TFR1 stage). 2. Percentage of patients who remain in DMR at the end of the consolidation period (week 48 of TFR1 stage), at 96 weeks and at 144 weeks after the start of the consolidation period of TFR1 stage. 3.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Percentage of patients in full treatment-free remission 144 weeks after the start of consolidation (end of the TFR1 stage.)

2.2 Background & Rationale

This study is constituted of two stage: Treatment-Free Remission 1 (TFR1) stage and Treatment-Free Remission 2 (TFR2) stage.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE2, Status: ACTIVE_NOT_RECRUITING

Target N: 124

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

Not Provided

4.2 Exclusion Criteria

Not Provided

11. Study Identifiers & Governance

Primary ID: 332181817887867375

Registry Number: NCT03874858

Sponsor: Novartis

Study Title: A Study of Full Treatment-free Remission in Patients With Chronic Myeloid Leukemia

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____