

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	Tumor-agnostic Precision Immuno-oncology and Somatic Targeting Rational for You (TAPISTRY) Phase II Platform Trial
Short Title:	Tumor-agnostic Precision Immuno-oncology and Somatic Targeting Rational for You (TAPISTRY) Platform Study
Protocol Number:	3397838386935247
NCT Number:	NCT04589845
Sponsor Name:	Roche
Investigational Product:	trastuzumab emtansine
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. All Cohorts: Independent Review Committee (IRC)-assessed Objective Response Rate (ORR) Based on Confirmed Objective Response (OR) per Response Evaluation Criteria in Solid Tumors, Version 1.1 (RECIST v1.1): Confirmed objective response indicates ≥4 weeks after initial documentation of response.
Secondary Obj.:	1. All Cohorts: IRC-assessed Duration of Response (DOR) per RECIST v1.1 2. All Cohorts: IRC-assessed Clinical Benefit Rate (CBR) per RECIST v1.1 3. All Cohorts: IRC-assessed Progression-free Survival (PFS) per RECIST v1.1

2.2 Background & Rationale

TAPISTRY is a Phase II, global, multicenter, open-label, multi-cohort study designed to evaluate the safety and efficacy of targeted therapies or immunotherapy as single agents or in rational, specified combinations in participants with unresectable, locally advanced or metastatic solid tumors determined to harbor specific oncogenic genomic alterations or who are tumor mutational burden (TMB)-high as identified by a validated next-generation sequencing (NGS) assay. Participants with solid tumors will be treated with a drug or drug regimen tailored to their NGS assay results at screening. Participants will be assigned to the appropriate cohort based on their genetic alteration(s). Treatment will be assigned on the basis of relevant oncogenotype, will have cohort-specific inclusion/exclusion criteria, and, unless otherwise specified, will continue until disease progression, loss of clinical benefit, unacceptable toxicity, participant or physician decision to discontinue, or death, whichever occurs first.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE2, Status: RECRUITING
Target N:	920

4. Subject Selection (Eligibility Criteria)

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4.1 Inclusion Criteria

- * Histologically or cytologically confirmed diagnosis of advanced and unresectable or metastatic solid malignancy
- * Measurable disease as defined by RECIST v1.1, RANO, or INRC
- * Performance status as follows: Participants aged ≥ 18 years: Eastern Cooperative Oncology Group (ECOG) Performance Status 0-2; Participants aged 16 to < 18 years: Karnofsky score $\geq 50\%$; Participants aged < 16 years: Lansky score $\geq 50\%$
- * For participants aged ≥ 18 and < 18 years: adequate hematologic and end-organ function
- * Disease progression on prior treatment, or previously untreated disease with no available acceptable treatment
- * Adequate recovery from most recent systemic or local treatment for cancer
- * Life expectancy ≥ 8 weeks
- * Ability to comply with the study protocol, in the investigator's judgment
- * For female participants of childbearing potential: Negative serum pregnancy test ≥ 14 days prior to initiating study treatment, agreement to remain abstinent or use single or combined contraception methods that result in a failure rate of $< 1\%$ per year for the period defined in the cohort-specific inclusion criteria; and agreement to refrain from donating eggs during the same period
- * For male participants: Willingness to remain abstinent or use acceptable methods of contraception as defined in the cohort-specific inclusion criteria
- * In addition to the general inclusion criteria above, participants must meet all of the cohort-specific inclusion criteria for the respective cohort

4.2 Exclusion Criteria

- * Current participation or enrollment in another therapeutic clinical trial
- * Any anticancer treatment within 2 weeks or 5 half-lives prior to start of study treatment
- * Whole brain radiotherapy within 14 days prior to start of study treatment
- * Stereotactic radiosurgery within 7 days prior to start of study treatment
- * Pregnant or breastfeeding, or intending to become pregnant during the study
- * History of or concurrent serious medical condition or abnormality in clinical laboratory tests that, in the investigator's judgment, precludes the participant's safe participation in and completion of the study or confounds the ability to interpret data from the study
- * Incomplete recovery from any surgery prior to the start of study treatment that would interfere with the determination of safety or efficacy of study treatment
- * Significant cardiovascular disease, such as New York Heart Association cardiac disease (Class II or higher), myocardial infarction, or cerebrovascular accident within 3 months prior to enrollment, unstable arrhythmias, or unstable angina
- * History of another active cancer within 5 years prior to screening that may interfere with the determination of safety or efficacy of study treatment with respect to the qualifying solid tumor malignancy
- * In addition to the general exclusion criteria above, in order to be enrolled in a treatment cohort of the study, participants must not meet any of the cohort-specific exclusion criteria

11. Study Identifiers & Governance

Primary ID: 3397838386935247

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Registry Number: NCT04589845

Sponsor: Roche

Study Title: Tumor-agnostic Precision Immuno-oncology and Somatic Targeting Rational for You (TAPISTRY) Platform Study

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____