

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 2 Study to Evaluate the Pharmacokinetics, Efficacy, and Safety of Subcutaneous Pembrolizumab Coformulated With Hyaluronidase (MK-3475A) in Participants With Relapsed or Refractory Classical Hodgkin Lymphoma (rrcHL) or Relapsed or Refractory Primary Mediastinal Large B-cell Lymphoma (rrPMBCL)
Short Title:	A Study to Evaluate the Safety and Efficacy of Pembrolizumab (MK-3475) Coformulated With Berahyaluronidase Alfa (MK-3475A) in Participants With Relapsed or Refractory Classical Hodgkin Lymphoma (rrcHL) or Relapsed or Refractory Primary Mediastinal Large B-cell Lymphoma (rrPMBCL)(MK-3475A-F65)
Protocol Number:	403403033384798878
NCT Number:	NCT06504394
Sponsor Name:	Merck
Investigational Product:	hyaluronidase
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Objective Response Rate (ORR) per Lugano Classification Criteria as Assessed by Investigator: ORR is defined as the percentage of the participants who had complete response (CR) or partial response (PR) and will be evaluated using computed tomography (CT) and positron emission tomography (PET)-CT. Response was assessed based on the International Working Group Criteria: Lugano Classification (Cheson et al, Journal of Clinical Oncology, 2014). CR is complete metabolic (no/minimal fluorodeoxyglucose [FDG] uptake) and radiologic response (target lesions regress to ≤ 1.5 cm in longest transverse diameter of a lesion) and no new lesions. PR is partial metabolic (moderate/high FDG uptake) and radiologic response ($\geq 50\%$ decrease in sum of product diameters for multiple lesions of up to 6 target measurable nodes and extranodal sites, no increase in lesions, and spleen regressed by $\geq 50\%$ in length beyond normal). The percentage of participants who experience CR or PR as assessed by investigator will be presented. 2. Maximum Concentration (C_{max}) of Pembrolizumab After Administration of SC Pembrolizumab Coformulated with Hyaluronidase: Blood samples will be collected at designated timepoints for the determination of C_{max}. C_{max} is defined as the peak concentration of pembrolizumab after administration of SC pembrolizumab coformulated with hyaluronidase. 3. Lowest Plasma Concentration (C_{trough}) of Pembrolizumab After Administration of SC Pembrolizumab Coformulated with Hyaluronidase: Blood samples will be collected at designated timepoints for the determination of C_{trough}. C_{trough} is defined as the lowest concentration of
--------------------	---

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

pembrolizumab after administration of SC pembrolizumab coformulated with hyaluronidase.

Secondary Obj.:

1. Duration of Response (DOR) per Lugano Classification Criteria as Assessed by Investigator
2. Number of Participants with Antidrug Antibodies (ADA) Level of Pembrolizumab After Administration of SC Pembrolizumab Coformulated with Hyaluronidase
3. Maximum Concentration (Cmax) of Pembrolizumab After Administration of SC Pembrolizumab Coformulated with Hyaluronidase at Steady State

2.2 Background & Rationale

The primary purpose of the study is to assess the pharmacokinetics (PK) profile of pembrolizumab following subcutaneous (SC) injection of pembrolizumab coformulated with hyaluronidase, and to evaluate the objective response rate (ORR) of pembrolizumab (+) berahyaluronidase alfa SC in adult participants with Relapsed or Refractory Classical Hodgkin Lymphoma (rrcHL) or Relapsed or Refractory Primary Mediastinal Large B-cell Lymphoma (rrPMBCL). There is no formal hypothesis to be tested for this study.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE2, Status: RECRUITING

Target N: 60

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

The main inclusion criteria include but are not limited to the following:

- * Histologically confirmed diagnosis of classical Hodgkin lymphoma (cHL) or primary mediastinal B-cell lymphoma (PMBCL)
- * Radiographically measurable cHL or PMBCL disease assessed by investigator as per Lugano classification
- * Have a life expectancy of >3 months
- * Human immunodeficiency virus (HIV)-infected participants must have well controlled HIV on antiretroviral therapy (ART)
- * Participants who are hepatitis B surface antigen (HBsAg) positive are eligible if they have received hepatitis B virus (HBV) antiviral therapy for at least 4 weeks, and have undetectable HBV viral load before enrollment
- * Participants with history of hepatitis C virus (HCV) infection are eligible if they have completed curative antiviral therapy at least 4 weeks before enrollment and HCV viral load is undetectable at screening
- * Has an Eastern Cooperative Oncology Group (ECOG) performance status of 0 to 1 assessed within 7 days before or on the day of the first dose of study intervention

4.2 Exclusion Criteria

The main exclusion criteria include but are not limited to the following:

- * Has clinically significant (i.e., active) cardiovascular disease

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Has pericardial effusion or clinically significant pleural effusion
- * Has known additional malignancy that is progressing or has required active treatment within the past 2 years
- * Diagnosis of immunodeficiency or is receiving systemic steroid therapy or any other form of immunosuppressive therapy within 7 days prior to the first dose of study intervention
- * Received prior monoclonal antibody within 4 weeks prior to first dose of study intervention or has not recovered (i.e., Grade 1 or at baseline) from adverse events (AEs) due to agents administered more than 4 weeks earlier
- * Received prior therapy with an anti-programmed cell death 1 protein (anti-PD-1), anti-programmed cell death ligand 1 (anti-PD-L1), or anti-programmed cell death ligand 2 (anti-PD-L2) agent, or with an agent directed to another stimulatory or coinhibitory T-cell receptor
- * Received prior systemic anticancer therapy including investigational agents within 4 weeks before the first dose of study intervention
- * Received prior radiotherapy within 2 weeks of start of study intervention, or has radiation-related toxicities, requiring corticosteroids
- * Received a live or live-attenuated vaccine within 30 days before first dose of study intervention
- * Is receiving systemic antineoplastic chemotherapy, immunotherapy, or biological therapy not specified in this protocol
- * Has known active central nervous system (CNS) metastases and/or carcinomatous meningitis
- * Active autoimmune disease that has required systemic treatment in the past 2 years
- * History of (noninfectious) pneumonitis/interstitial lung disease that required steroids or has current pneumonitis/interstitial lung disease
- * Active infection requiring systemic therapy except certain protocol-specified therapies
- * Concurrent active hepatitis B and hepatitis C virus infection
- * Has undergone solid organ transplant at any time, or prior allogeneic hematopoietic stem cell transplant (SCT) within the last 5 years

11. Study Identifiers & Governance

Primary ID: 403403033384798878

Registry Number: NCT06504394

Sponsor: Merck

Study Title: A Study to Evaluate the Safety and Efficacy of Pembrolizumab (MK-3475) Coformulated With Berahyaluronidase Alfa (MK-3475A) in Participants With Relapsed or Refractory Classical Hodgkin Lymphoma (rrcHL) or Relapsed or Refractory Primary Mediastinal Large B-cell Lymphoma (rrPMBCL)(MK-3475A-F65)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____