

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase II, Randomized, Double-Blind, Placebo-Controlled, Multicenter Study to Evaluate the Efficacy, Safety, and Pharmacokinetics of Obinutuzumab in Adolescent Patients With Active Class III or IV Lupus Nephritis, Including an Evaluation of Open Label Safety and PK in a Cohort of Pediatric Patients (Aged 5 to < 12)
Short Title:	A Study to Evaluate the Efficacy, Safety, and Pharmacokinetics of Obinutuzumab in Adolescents With Active Class III or IV Lupus Nephritis and the Safety and PK of Obinutuzumab in Pediatric Participants
Protocol Number:	488634691547466382
NCT Number:	NCT05039619
Sponsor Name:	Roche
Investigational Product:	paracetamol
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Percentage of Participants who Achieve a Complete Renal Response (CRR) (AP): CRR is defined as achievement of all of the following: * Urinary protein-to-creatinine ratio (UPCR) ≤ 0.5 g/g * Estimated Glomerular Filtration Rate (eGFR) $\geq 85\%$ of baseline * No occurrence of intercurrent events 2. Percentage of Participants with Adverse Events (PP)
Secondary Obj.:	1. Percentage of Participants Achieving a CRR (AP) 2. Percentage of Participants who Achieve CRR with Successful Prednisone Taper (AP) 3. Percentage of Participants who Achieve a PRR (AP)

2.2 Background & Rationale

This phase II, randomized, double-blind, placebo-controlled study is designed to evaluate the safety, efficacy and pharmacokinetics (PK) of obinutuzumab in adolescent participants (AP) aged 12 to less than 18 with biopsy-confirmed proliferative lupus nephritis (LN). It will also evaluate open label safety and PK of obinutuzumab in pediatric participants (PP), aged 5 to ≤ 12 with LN.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE2, Status: RECRUITING
Target N:	40

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Participants who are age 12 to <18 years at the time of randomization
- * Participants who are age 5 to <12 years (younger participant cohort) at the time of randomization once recruitment is open. (Investigators will be notified by the Sponsor when recruitment is open to this younger population)
- * International Society of Nephrology and the Renal Pathology Society (ISN/RPS) 2003 Class III or IV active LN demonstrated on renal biopsy performed in the 12 months prior to or during screening
- * Class V disease may be present in addition to Class III or IV LN, but participants with isolated Class V disease are not eligible
- * Diagnosis of SLE according to the Systemic Lupus International Collaborating Clinics (SLICC) 2012 criteria
- * Significant proteinuria defined by a UPCR above > 0.5 based on a first-morning void (FMV) collection at screening
- * During the 12 months prior to or during screening, all participants must have received at least one dose of pulse-range IV methylprednisolone (typically 30 mg/kg, maximum of 1000 mg per dose) or equivalent for the treatment of the current episode of active LN.

4.2 Exclusion Criteria

- * Severe, active central nervous system (CNS) SLE, including retinitis, poorly controlled seizure disorder, acute confusional state, myelitis, stroke, cerebellar ataxia, or dementia
- * Sclerosis in >50% of glomeruli on renal biopsy
- * Purely chronic Class III(c) or Class IV(c) disease on renal biopsy, defined as the absence of any active lesions
- * Presence of rapidly progressive glomerulonephritis
- * Pure Class V LN
- * Intolerance or contraindication to study therapies
- * Active infection of any kind (excluding fungal infection of nail beds) or any major episode of infection requiring hospitalization or treatment with IV anti-infective medications within 4 weeks prior to screening, or completion of oral anti-infectives within 2 weeks prior to randomization
- * History of or currently active primary or secondary immunodeficiency, including known history of HIV infection and other severe Immunodeficiency blood disorders
- * History of serious recurrent or chronic infection
- * History of or current cancer, including solid tumors, hematological malignancies, and carcinoma in situ (except basal cell carcinoma and squamous cell carcinoma of the skin that have been excised and cured) within the past 5 years
- * Significant or uncontrolled concomitant medical disease which, in the investigator's opinion, would preclude participant participation
- * Currently active alcohol or drug abuse or history of alcohol or drug abuse

11. Study Identifiers & Governance

Primary ID: 488634691547466382

Registry Number: NCT05039619

Sponsor: Roche

Study Title: A Study to Evaluate the Efficacy, Safety, and Pharmacokinetics of Obinutuzumab in

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Adolescents With Active Class III or IV Lupus Nephritis and the Safety and PK of Obinutuzumab in Pediatric Participants

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____