

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	KEYMAKER-U01 Substudy 01H: A Phase 2, Randomized, Umbrella Study With Rolling Arms of Investigational Agents in Participants With Previously Treated Stage IV Nonsquamous Non-small Cell Lung Cancer (NSCLC)
Short Title:	A Study of Investigational Agents in Participants With Previously Treated Stage IV Nonsquamous Non-small Cell Lung Cancer (NSCLC) (MK-3475-01H/KEYMAKER-U01)
Protocol Number:	558163847581215514
NCT Number:	NCT06780085
Sponsor Name:	Merck
Investigational Product:	docetaxel
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Objective Response Rate (ORR): ORR is defined as the percentage of participants with Complete Response (CR: Disappearance of all target lesions) or a Partial Response (PR: At least a 30% decrease in the sum of diameters of target lesions) per Response Evaluation Criteria In Solid Tumors Version 1.1 (RECIST 1.1). The percentage of participants who experience CR or PR as assessed by Blinded Independent Central Review (BICR) will be presented. 2. Percentage of Participants with at Least One Adverse Event (AE): An AE is defined as any unfavorable and unintended sign, symptom, disease, or worsening of preexisting condition temporally associated with study treatment and irrespective of causality to study treatment. The percentage of participants who experience an AE will be reported. 3. Percentage of Participants Who Discontinued Medication Due to an AE: An AE is defined as any unfavorable and unintended sign, symptom, disease, or worsening of preexisting condition temporally associated with study treatment and irrespective of causality to study treatment. The percentage of participants who discontinue study intervention due to an AE will be reported.
Secondary Obj.:	1. Duration of Response (DOR) 2. Progression-free Survival (PFS) 3. Overall Survival (OS)

2.2 Background & Rationale

Researchers are looking for new ways to treat metastatic nonsquamous non-small cell lung cancer (NSCLC) that has been treated before. Metastatic means the cancer has spread to other parts of the body. Nonsquamous means the cancer did not start in squamous cells, which are flat cells that line the inside of the lungs.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE2, Status: RECRUITING

Target N: 96

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

The main inclusion criteria include but are not limited to the following:

- * Histologically or cytologically confirmed diagnosis of Stage IV nonsquamous non-small cell lung cancer (NSCLC)
- * Documented disease progression per RECIST 1.1 after receiving an anti-programmed cell death 1 protein (PD-1)/programmed cell death ligand 1 (PD-L1) treatment and platinum-based chemotherapy
- * Confirmation per local test report that epidermal growth factor receptor negative (EGFR-), anaplastic lymphoma kinase negative (ALK-), or c ros oncogene 1 negative (ROS1-) directed therapy is not indicated as primary therapy
- * Measurable disease per RECIST 1.1 as assessed by investigator and verified by BICR
- * Life expectancy of at least 3 months
- * An Eastern Cooperative Oncology Group (ECOG) performance status of 0 to 1 assessed within 7 days before randomization
- * Is an individual of any sex/gender who is at least 18 years of age at the time of providing the informed consent
- * Has adequate organ function
- * If capable of producing sperm refrains from donating sperm plus either abstains from penile-vaginal intercourse or uses a penile/external condom, with contraceptive use consistent with local regulations
- * Participant/participants of childbearing potential (POCBP) is not pregnant and has a negative highly sensitive pregnancy test; and is not breastfeeding and uses a highly effective contraceptive method
- * Archival tumor tissue sample of a tumor lesion not previously irradiated has been provided
- * Has provided tissue prior to treatment randomization from a newly obtained formalin-fixed sample from a new biopsy
- * Human Immunodeficiency Virus (HIV)-infected participants must have well controlled HIV on antiretroviral therapy (ART)
- * Participants who are hepatitis B surface antigen (HBsAg) positive have received hepatitis B virus (HBV) antiviral therapy for at least 4 weeks and have undetectable HBV viral load prior to randomization
- * Participants with history of hepatitis C virus (HCV) infection are eligible if HCV viral load is undetectable at screening

4.2 Exclusion Criteria

The main exclusion criteria include but are not limited to the following:

- * Diagnosis of small cell lung cancer or, for mixed tumors, presence of small cell elements
- * Received radiation therapy to the lung
- * Has uncontrolled or significant cardiovascular disorder prior to randomization
- * Has clinically severe pulmonary compromise resulting from intercurrent pulmonary illnesses
- * Participants who have adverse events (AEs) (other than alopecia) due to previous anticancer therapies must have recovered to ?Grade 1 or baseline

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Has known severe hypersensitivity (?Grade 3) to study intervention and/or any of its excipients
- * Has evidence of ongoing uncontrolled systemic bacterial, fungal, or viral infection (including HIV infection)
- * Has clinically significant corneal disease
- * Has received prior radiotherapy within 2 weeks of start of study intervention, or radiation related toxicities, requiring corticosteroids
- * Has inadequate washout period prior to randomization
- * Has received a live or live-attenuated vaccine within 30 days before the first dose of study intervention
- * Has diagnosis of immunodeficiency or is receiving chronic systemic steroid therapy
- * Has received an investigational agent or has used an investigational device within 4 weeks prior to study intervention administration
- * Has known additional malignancy that is progressing or has required active treatment within the past 3 years
- * Has known untreated central nervous system (CNS) metastases and/or carcinomatous meningitis
- * Has evidence of any leptomeningeal disease
- * Has history of (noninfectious) pneumonitis/ interstitial lung disease (ILD) that required steroids or has current pneumonitis/ILD, and/or suspected ILD/pneumonitis
- * Has active autoimmune disease that has required systemic treatment in the past 2 years
- * Has active infection requiring systemic therapy
- * HIV-infected participants with a history of Kaposi's sarcoma and/or Multicentric Castleman's Disease
- * Has active inflammatory bowel disease requiring immunosuppressive medication or previous clear history of inflammatory bowel disease
- * Has known history of, or active, neurologic paraneoplastic syndrome
- * Has history of allogeneic tissue/solid organ transplant
- * Have not adequately recovered from major surgery or have ongoing surgical complications

11. Study Identifiers & Governance

Primary ID: 558163847581215514

Registry Number: NCT06780085

Sponsor: Merck

Study Title: A Study of Investigational Agents in Participants With Previously Treated Stage IV Nonsquamous Non-small Cell Lung Cancer (NSCLC) (MK-3475-01H/KEYMAKER-U01)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____