

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase II, Multicenter, Randomized, Open-label Study to Evaluate the Safety and Efficacy of 400 mg of Ribociclib in Combination With Non-steroidal Aromatase Inhibitors for the Treatment of Pre- and Postmenopausal Women With Hormone Receptor-positive, HER2-negative Advanced Breast Cancer Who Received no Prior Therapy for Advanced Disease
Short Title:	Study of 2 Ribociclib Doses in Combination With Aromatase Inhibitors in Women With HR+, HER2- Advanced Breast Cancer
Protocol Number:	726292173838059615
NCT Number:	NCT03822468
Sponsor Name:	Novartis
Investigational Product:	goserelin
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Overall Response Rate (ORR): ORR defined as the percentage of participants with best overall response (BOR) of confirmed complete response (CR) or partial response (PR) assessed by local investigators according to RECIST 1.1. CR: Disappearance of all lesions with lymph nodes measuring ≤ 10 mm. PR: At least a 30% decrease in the sum of diameter of all target lesions, taking as reference the baseline sum of diameters.
Secondary Obj.:	1. Change From Baseline in QTc (With Fridericia's Correction) at Cycle 1 Day 15 (at 2 Hours Post-dose) 2. Progression-free Survival (PFS) 3. Clinical Benefit Rate (CBR)

2.2 Background & Rationale

The purpose of the study was to evaluate the safety and efficacy of a reduced ribociclib starting dose of 400 mg in combination with a non-steroidal aromatase inhibitor (NSAI) (letrozole or anastrozole) for the treatment of pre- and postmenopausal women with hormone receptor-positive (HR-positive), HER2-negative advanced breast cancer (aBC) who have received no prior therapy for advanced disease. Premenopausal women were required to receive goserelin in both treatment arms.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE2, Status: COMPLETED
Target N:	376

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

Not Provided

4.2 Exclusion Criteria

- * Patient with symptomatic visceral disease or any disease burden that makes the patient ineligible for endocrine therapy per the investigator's judgment.
- * Patient who received any prior systemic anti-cancer therapy(including endocrine therapy, chemotherapy, prior CDK4/6 inhibitors) for aBC. Patients who received neo-/adjuvant therapy for breast cancer are eligible.
- * Patient is concurrently using other anti-cancer therapy.
- * Patient has had major surgery within 14 days prior to starting study drug or has not recovered from major toxicities.
- * Patient has received extended-field radiotherapy ? 4 weeks or limited field radiotherapy ? 2 weeks prior to randomization, and has not recovered to grade 1 or better from related side effects of such therapy (with the exception of alopecia or other toxicities not considered a safety risk for the patient at investigator's discretion). Patients in whom ?
- * 25% of the bone marrow has been previously irradiated are also excluded.
- * Patient has a concurrent malignancy or malignancy within 3 years of the randomization date, with the exception of adequately treated basal or squamous cell skin carcinoma, or curatively resected cervical carcinoma in situ.
- * Patients with central nervous system (CNS) involvement unless they meet specific stability criteria.
- * Patient has clinically significant, uncontrolled heart disease and/or cardiac repolarization abnormality.
- * Patient is currently receiving or has received systemic corticosteroids ? 2 weeks prior to starting study drug, and has not fully recovered from side effects of such treatment.

Other protocol-defined Inclusion/Exclusion may apply.

11. Study Identifiers & Governance

Primary ID: 726292173838059615

Registry Number: NCT03822468

Sponsor: Novartis

Study Title: Study of 2 Ribociclib Doses in Combination With Aromatase Inhibitors in Women With HR+, HER2- Advanced Breast Cancer

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____