

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	An Open-Label Study to Investigate the Safety, Tolerability, and Pharmacokinetics/Pharmacodynamics of Risdiplam (RO7034067) in Adult and Pediatric Patients With Spinal Muscular Atrophy
Short Title:	A Study of Risdiplam (RO7034067) in Adult and Pediatric Participants With Spinal Muscular Atrophy
Protocol Number:	730147882578611099
NCT Number:	NCT03032172
Sponsor Name:	Roche
Investigational Product:	risdiplam
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Number of Participants With Adverse Events (AEs): An AE was any untoward medical occurrence in a clinical investigation participant administered a pharmaceutical product, regardless of causal attribution. An AE can therefore be any unfavorable and unintended sign (including an abnormal laboratory finding), symptom, or disease temporally associated with the use of a medicinal product, whether or not considered related to the medicinal product. 2. Number of Participants Who Discontinued Treatment Due to AEs: An AE was any untoward medical occurrence in a clinical investigation participant administered a pharmaceutical product, regardless of causal attribution. An AE can therefore be any unfavorable and unintended sign (including an abnormal laboratory finding), symptom, or disease temporally associated with the use of a medicinal product, whether or not considered related to the medicinal product. Participants who discontinued treatment due to AEs are reported here. 3. Number of Participants With Shift in Puberty Status From Baseline: Tanner staging of sexual development is a scale used to assess physical maturation as children transition through adolescence into adulthood. Scale defines physical development based on the following characteristics: pubic hair, penis, and testes development in boys; and pubic hair and breast development in girls. It consists of 5 stages, Stage I (prepubertal) to Stage V (mature adult). Participants under 9 years at screening began Tanner staging assessments at 1st visit following their 9th birthday. Tanner data are presented in three categories: Normal (expected stage of puberty based on participant's age at post-baseline visit), Delayed (pubertal development is behind expectations for age at post-baseline visit), & Missing (participant did not attend scheduled visit). Tanner staging assessments were scheduled for participants aged 9-17 years but were also
--------------------	--

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

conducted in some older participants, up to age 22. Shift in puberty status from baseline to each week has been represented here.

Secondary Obj.: 1. Survival of Motor Neuron (SMN) Protein Levels in Blood

2.2 Background & Rationale

This is a multi-center, exploratory, non-comparative, and open-label study to investigate the safety, tolerability, PK, and PK/PD relationship of risdiplam in adults, children and infants with Spinal Muscular Atrophy (SMA) previously enrolled in Study BP29420 (Moonfish) with the splicing modifier RO6885247 or previously treated with nusinersen, olesoxime or AVXS-101.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE2, Status: COMPLETED

Target N: 174

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Confirmed diagnosis of 5q-autosomal recessive SMA
- * Previous enrollment in Study BP29420 (Moonfish) with the splicing modifier RO6885247 or previous treatment with any of the following: 1.) Nusinersen (defined as having received ≥ 4 doses of nusinersen, provided that the last dose was received ≥ 90 days prior to screening) or 2.) Olesoxime (provided that the last dose was received ≤ 12 months and ≥ 90 days prior to screening) or 3.) AVXS-101 (provided that the time of treatment was ≥ 12 months prior to screening)
- * Adequately recovered from any acute illness at the time of screening and considered well enough to participate in the opinion of the Investigator
- * For women of childbearing potential: negative blood pregnancy test at screening, agreement to remain abstinent (refrain from heterosexual intercourse) or use contraceptive measures, and agreement to refrain from donating eggs for at least 28 days after the final dose of study drug
- * For men: agreement to remain abstinent (refrain from heterosexual intercourse) or use contraceptive measures and agreement to refrain from donating sperm
- * For participants aged 2 years or younger at screening: 1.) Parent or caregiver of participant is willing to consider nasogastric, naso-jejunal or gastrostomy tube placement, as recommended by the Investigator, during the study to maintain safe hydration, nutrition and treatment delivery; 2.) Parent or caregiver of participant is willing to consider the use of non-invasive ventilation, as recommended by the Investigator during the study

4.2 Exclusion Criteria

- * Inability to meet study requirements
- * Concomitant participation in any investigational drug or device study
- * Previous participation in any investigational drug or device study within 90 days prior to screening, or 5 half-lives of the drug, whichever is longer with the exception of studies of olesoxime, AVXS-101, or nusinersen
- * Any history of gene or cell therapy, with the exception of AVXS-101

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Unstable gastrointestinal, renal, hepatic, endocrine, or cardiovascular system diseases as considered to be clinically significant by the Investigator
- * Inadequate venous or capillary blood access for the study procedures, in the opinion of the Investigator
- * For patients aged ≤ 2 years, hospitalization for a pulmonary event within 2 months prior to screening and pulmonary function not fully recovered at the time of screening
- * Lactating women
- * Suspicion of regular consumption of drugs of abuse
- * For adults and adolescents only, positive urine test for drugs of abuse or alcohol at screening or Day -1 visit
- * Presence of clinically significant electrocardiogram (ECG) abnormalities before study drug administration from average of triplicate measurement or cardiovascular disease
- * History of malignancy if not considered cured
- * For participants aged ≥ 6 years, significant risk for suicidal behavior, in the opinion of the Investigator as assessed by the Columbia-Suicide Severity Rating Scale (C-SSRS)
- * Any major illness within one month before the screening examination or any febrile illness within one week prior to screening and up to first dose administration
- * Recently initiated treatment for spinal muscular atrophy (within ≤ 6 weeks prior to enrollment) with oral salbutamol or another beta 2-adrenergic agonist taken orally
- * Any prior use of chloroquine, hydroxychloroquine, retigabine, vigabatrin or thioridazine, is not allowed
- * Ascertained or presumptive hypersensitivity (e.g., anaphylactic reaction) to risdiplam or to the constituents of its formulation
- * Concomitant disease or condition that could interfere with, or treatment of which might interfere with, the conduct of the study, or that would, in the opinion of the Investigator, pose an unacceptable risk to the participant in this study
- * Recent history (less than one year) of ophthalmological diseases
- * Any prior use of an inhibitor or inducer of CYP1A2 or CYP3A4 taken within 2 weeks (or within 5 elimination half-lives, whichever is longer) prior to dosing

11. Study Identifiers & Governance

Primary ID: 730147882578611099

Registry Number: NCT03032172

Sponsor: Roche

Study Title: A Study of Risdiplam (RO7034067) in Adult and Pediatric Participants With Spinal Muscular Atrophy

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____