

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	Open Label Phase 2 Study of Tisotumab Vedotin for Locally Advanced or Metastatic Disease in Solid Tumors
Short Title:	Efficacy and Safety Study of Tisotumab Vedotin for Patients With Solid Tumors
Protocol Number:	766342340119063828
NCT Number:	NCT03485209
Sponsor Name:	Pfizer
Investigational Product:	cisplatin
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Confirmed Objective Response Rate (ORR) (Parts A, B, C, D, E, F, and G): Proportion of patients who achieve a confirmed complete response (CR) or partial response (PR) according to RECIST v1.1 as assessed by the investigator
Secondary Obj.:	1. Incidence of Adverse Events (AEs) 2. Confirmed and Unconfirmed ORR 3. Disease Control Rate (DCR)

2.2 Background & Rationale

This trial will study tisotumab vedotin to find out whether it is an effective treatment alone or with other anticancer drugs for certain solid tumors and what side effects (unwanted effects) may occur. There are seven parts to this study.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE2, Status: ACTIVE_NOT_RECRUITING
Target N:	352

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Parts A, B, and C
- * Relapsed, locally-advanced or metastatic colorectal or pancreatic cancer, sqNSCLC, or HNSCC participants who are not candidates for standard therapy.
- * All participants must have experienced disease progression on or after their most recent systemic therapy.
- * Colorectal cancer (closed to enrollment): participants must have received prior therapy with each of following agents, if eligible: a fluoropyrimidine, oxaliplatin, irinotecan, and/or bevacizumab. Participants should have received no more than 3 systemic regimens in the metastatic setting.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

* sqNSCLC (closed to enrollment): Participants with NSCLC must have predominant squamous histology. Participants must have received prior therapy with a platinum-based treatment and a checkpoint inhibitor (CPI), if eligible. Participants should have received no more than 3 lines of systemic therapy in the metastatic setting.

* Participants eligible for a tyrosine kinase inhibitor should have received such therapy. These participants should have received no more than 4 lines of systemic therapy in the metastatic setting.

* Exocrine pancreatic adenocarcinoma (closed to enrollment): Participants with exocrine pancreatic adenocarcinoma must have predominant adenocarcinoma histology. Participants must have received prior therapy with a gemcitabine-based or 5FU-based regimen, if eligible, and should have received no more than 1 systemic regimen in the unresectable or metastatic setting.

* HNSCC (closed to enrollment): Participants with HNSCC in Part C must have received prior therapy with a platinum-based regimen and/or a checkpoint inhibitor (CPI), if eligible, and must have experienced disease progression following such therapy. Participants should have received no more than 3 systemic lines of therapy in the recurrent or metastatic setting.

* Part E

* Participants with HNSCC must have experienced disease progression on or after their most recent systemic therapy. Participants should have received no more than 1 or 2 systemic lines of therapy in the recurrent/metastatic setting as specified below. Participants must have received a platinum-based regimen and a PD-(L)1 inhibitor.

* Parts D, F, and G

* Part D is closed to enrollment. Part F and Part G will enroll only participants with HNSCC.

* Participants with HNSCC must have received no previous systemic therapy in the recurrent or metastatic disease setting.

* Part D only

* Participants with NSCLC must have histologically or cytologically documented squamous cell NSCLC and must have received no previous systemic therapy for metastatic disease or radiation therapy to the lung that is ≥ 30 Gy within 6 months of the first dose of study treatment.

* PD-L1 biomarker expression as determined by a PD-L1 IHC assay should be available

* Part F only

* Participants must have CPS ≥ 1 by local PD-L1 IHC assay to be eligible for enrollment. Participants must be able to submit a tissue sample for retrospective PD-L1 testing. Tissue may be fresh biopsy or archival, collected within 2 years of Cycle 1 Day 1.

* Part G only

* Non-EU eligibility criteria: No CPS requirement for the cohort evaluating tisotumab vedotin in combination with pembrolizumab and carboplatin.

* EU-specific eligibility criteria: Participants must have a CPS ≥ 1 by local PD-L1 IHC assay.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Participants must be able to submit a tissue sample for retrospective PD-L1 testing. Tissue may be fresh biopsy or archival, collected within 2 years of Cycle 1 Day 1.
- * Baseline measurable disease as measured by RECIST v1. 1.
- * Eastern Cooperative Oncology Group (ECOG) Performance Status score of 0 or 1.

4.2 Exclusion Criteria

- * Participants with primary neuroendocrine or sarcomatoid histologies. For HNSCC, participants may not have a primary site of nasopharynx or salivary gland.
- * Active bleeding conditions
- * Ocular surface disease at the time of enrollment (Note: cataract is not considered active ocular surface disease for this protocol)
- * Other cancer: known past or current malignancy other than inclusion diagnosis.
- * Uncontrolled tumor-related pain
- * Inflammatory lung disease. Participants with pulmonary disease are allowed if systemic steroids and long-term oxygen are not required
- * Peripheral neuropathy greater than or equal to Grade 2
- * Active brain metastasis
- * Ongoing clinically significant toxicity associated with prior treatment (including radiotherapy or surgery).
- * Part D, F, and G Only: Prior therapy with an anti-PD-1, anti-PD-L1, or anti-PD-L2 agent or with an agent directed to another stimulatory or co-inhibitory T-cell receptor.

11. Study Identifiers & Governance

Primary ID: 766342340119063828
Registry Number: NCT03485209
Sponsor: Pfizer
Study Title: Efficacy and Safety Study of Tisotumab Vedotin for Patients With Solid Tumors

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____