

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase II, Open-label Study to Investigate the Pharmacokinetics and Safety of Risdiplam in Infants With Spinal Muscular Atrophy
Short Title:	A Study to Investigate the Pharmacokinetics and Safety of Risdiplam in Infants With Spinal Muscular Atrophy
Protocol Number:	849699474298655348
NCT Number:	NCT05808764
Sponsor Name:	Roche
Investigational Product:	risdiplam
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Plasma Concentrations of Risdiplam 2. Area Under the Plasma Concentration-Time Curve (AUC) of Risdiplam 3. Steady-state Concentration (Css) of Risdiplam
Secondary Obj.:	Not Provided

2.2 Background & Rationale

This study will evaluate the pharmacokinetics (PK) and safety of risdiplam in participants with spinal muscular atrophy (SMA) under 20 days of age at first dose.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE2, Status: RECRUITING
Target N:	10

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Male or female newborn infant aged ≤ 20 days at first dose
- * Newborn infants with genetic diagnosis of 5q-autosomal recessive SMA or newborn infants identified as positive for SMA via newborn screening or via prenatal testing.
- * Gestational age equal to or greater than 37 weeks
- * Receiving adequate nutrition and hydration at the time of screening
- * Adequately recovered from any acute illness at baseline and considered well enough to participate in the study
- * Parent/caregiver is willing to consider nasogastric, nasojejunal, or gastrostomy tube placement during the study to maintain safe hydration, nutrition, and treatment delivery, if recommended by the investigator.

4.2 Exclusion Criteria

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Presence of clinical symptoms or signs consistent with SMA Type 0
- * In the opinion of the investigator, inadequate venous or capillary blood access for the study procedures
- * Systolic blood pressure or diastolic blood pressure or heart rate abnormalities
- * Presence of clinically relevant electrocardiogram (ECG) abnormalities
- * The infant (or the person breastfeeding the infant) taking any of the following: any inhibitor of CYP3A4 taken within 2 weeks (or within 5 times the elimination half-life, whichever is longer) prior to dosing, any inducer of CYP3A4 taken within 4 weeks (or within 5 times the elimination half-life, whichever is longer prior to dosing, and/or use of any multidrug and toxin extrusion (MATE) substrates taken within 2 weeks (or within 5 times the elimination half-life, whichever is longer) prior to dosing
- * Concurrent or previous administration of nusinersen or onasemnogene abeparvovec
- * Clinically significant abnormalities in laboratory test

11. Study Identifiers & Governance

Primary ID: 849699474298655348

Registry Number: NCT05808764

Sponsor: Roche

Study Title: A Study to Investigate the Pharmacokinetics and Safety of Risdiplam in Infants With Spinal Muscular Atrophy

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____