

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase II, Open-Label, Multicenter, Platform Study Evaluating the Efficacy and Safety of Biomarker-Driven Therapies in Patients With Persistent or Recurrent Rare Epithelial Ovarian Tumors
Short Title:	A Study Evaluating the Efficacy and Safety of Biomarker-Driven Therapies in Patients With Persistent or Recurrent Rare Epithelial Ovarian Tumors
Protocol Number:	897395762950279385
NCT Number:	NCT04931342
Sponsor Name:	Roche
Investigational Product:	cyclophosphamide
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Confirmed Objective Response Rate (ORR): Confirmed ORR is defined as the proportion of participants with a confirmed complete response (CR) or partial response (PR) (demonstrated on two consecutive occasions ≥ 4 weeks apart), as determined by the investigator according to RECIST v1.1.
Secondary Obj.:	1. Duration of Response (DOR) 2. Disease Control Rate (DCR) 3. Progression Free Survival (PFS)

2.2 Background & Rationale

This study will evaluate the efficacy and safety of multiple biomarker-selected treatments in patients with persistent or recurrent rare epithelial ovarian, fallopian tube, or primary peritoneal tumors. Enrollment will take place in two phases: a preliminary phase followed by a potential expansion phase.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE2, Status: ACTIVE_NOT_RECRUITING
Target N:	176

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

* Persistent or recurrent EOC that meets the following criteria: Histologically confirmed non-high-grade serous, non-high-grade endometrioid epithelial ovarian, fallopian tube, or primary peritoneal cancer, including but not limited to low-grade serous ovarian carcinoma, clear cell carcinoma, mucinous carcinoma, carcinosarcoma, undifferentiated carcinoma, seromucinous carcinoma, malignant Brenner tumors, Grades 1 or 2 endometrioid carcinoma,

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

mesonephric-like adenocarcinoma and small cell carcinoma of the ovary, hypercalcemic type (SCCOHT). Disease that is not amenable to curative surgery

- * Measurable disease (at least one target lesion) according to RECIST v1.1
- * Previous treatment with one to four lines of therapy, at least one of which was platinum-based. Hormonal therapy does not count as a line of therapy.
- * Platinum-resistant disease, defined as disease progression during or within 6 months of last platinum therapy, with the following exception: Participants with primary platinum-refractory disease are excluded.
- * Submission of a representative tumor specimen that is suitable for next-generation sequencing (NGS) testing and estrogen receptor immunohistochemistry (ER IHC) to determine treatment arm assignment and for central pathology review.
- * Submission of the local pathology report and, if available, any associated stained slides that supported the local diagnosis of the histology (to be used for central pathology review)
- * Eastern Cooperative Oncology Group (ECOG) Performance Status of 0 or 1
- * Adequate hematologic and end-organ function
- * For women of childbearing potential: agreement to remain abstinent or use contraception, and agreement to refrain from donating eggs (if applicable)
- * In addition to the general inclusion criteria above, participants must meet all of the arm-specific inclusion criteria for the respective arm

General Exclusion Criteria:

- * Pregnant or breastfeeding, or intending to become pregnant or breastfeed during the study
- * Primary platinum-refractory disease, defined as progression during or within 4 weeks after the last dose of the first-line platinum treatment
- * Histologic diagnosis of high-grade serous or high-grade endometrioid ovarian, fallopian tube, or primary peritoneal cancer
- * Current diagnosis of solely borderline epithelial ovarian tumor
- * Current diagnosis of non-epithelial ovarian tumors
- * Current diagnosis of synchronous primary endometrial cancer
- * Prior history of primary endometrial cancer, with the following exception: a prior diagnosis of primary endometrial cancer is permitted if it meets all of the following conditions: Stage IA, no lymphovascular invasion, International Federation of Gynecology and Obstetrics Grade 1 or 2, not a high-grade subtype.
- * Uncontrolled pleural effusion, pericardial effusion, or ascites requiring recurrent drainage procedures
- * Symptomatic, untreated, or actively progressing CNS metastases
- * Severe infection within 4 weeks prior to initiation of study treatment
- * Treatment with chemotherapy, radiotherapy, antibody therapy or other immunotherapy, gene therapy, vaccine therapy, or investigational therapy within 28 days prior to initiation of study treatment
- * Treatment with hormonal therapy within 14 days prior to initiation of study treatment
- * In addition to the general exclusion criteria above, participants can not meet any of the arm-specific exclusion criteria for the respective arm

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4.2 Exclusion Criteria

- * Pregnant or breastfeeding, or intending to become pregnant or breastfeed during the study
- * Primary platinum-refractory disease, defined as progression during or within 4 weeks after the last dose of the first-line platinum treatment
- * Histologic diagnosis of high-grade serous or high-grade endometrioid ovarian, fallopian tube, or primary peritoneal cancer
- * Current diagnosis of solely borderline epithelial ovarian tumor
- * Current diagnosis of non-epithelial ovarian tumors
- * Current diagnosis of synchronous primary endometrial cancer
- * Prior history of primary endometrial cancer, with the following exception: a prior diagnosis of primary endometrial cancer is permitted if it meets all of the following conditions: Stage IA, no lymphovascular invasion, International Federation of Gynecology and Obstetrics Grade 1 or 2, not a high-grade subtype.
- * Uncontrolled pleural effusion, pericardial effusion, or ascites requiring recurrent drainage procedures
- * Symptomatic, untreated, or actively progressing CNS metastases
- * Severe infection within 4 weeks prior to initiation of study treatment
- * Treatment with chemotherapy, radiotherapy, antibody therapy or other immunotherapy, gene therapy, vaccine therapy, or investigational therapy within 28 days prior to initiation of study treatment
- * Treatment with hormonal therapy within 14 days prior to initiation of study treatment
- * In addition to the general exclusion criteria above, participants can not meet any of the arm-specific exclusion criteria for the respective arm

11. Study Identifiers & Governance

Primary ID:	897395762950279385
Registry Number:	NCT04931342
Sponsor:	Roche
Study Title:	A Study Evaluating the Efficacy and Safety of Biomarker-Driven Therapies in Patients With Persistent or Recurrent Rare Epithelial Ovarian Tumors

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____