

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 1b/2 Study of BMS-813160 in Combination With Chemotherapy or Nivolumab in Patients With Advanced Solid Tumors
Short Title:	A Study of BMS-813160 in Combination With Chemotherapy or Nivolumab in Participants With Advanced Solid Tumors
Protocol Number:	22720114948792005
NCT Number:	NCT03184870
Sponsor Name:	Bristol Myers Squibb
Investigational Product:	fluorouracil
Phase:	PHASE1, PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Number of Participants Experiencing Adverse Events (AEs): An Adverse Event (AE) is defined as any new untoward medical occurrence or worsening of a pre-existing medical condition in a clinical investigation participant administered study treatment and that does not necessarily have a causal relationship with this treatment. Adverse events are graded on a scale from 1 to 5, with Grade 1 being mild and asymptomatic; Grade 2 is moderate requiring minimal, local or noninvasive intervention; Grade 3 is severe or medically significant but not immediately life-threatening; Grade 4 events are usually severe enough to require hospitalization; Grade 5 events are fatal. 2. Number of Participants Experiencing Serious Adverse Events (SAEs): A Serious Adverse Event (SAE) is defined as any untoward medical occurrence that, at any dose results in death, is life-threatening (defined as an event in which the participant was at risk of death at the time of the event; it does not refer to an event which hypothetically might have caused death if it were more severe), requires inpatient hospitalization or causes prolongation of existing hospitalization. 3. Number of Participants Experiencing Dose-Limiting Toxicities (DLTs): Dose-limiting toxicities (DLTs) are severe adverse effects (AEs) that are attributed to BMS-813160 or the combination regimen and define the maximum tolerated dose of a medicine. DLTs will be defined based on the incidence, duration and grade of AEs for which no alternate cause can be identified. AEs will be evaluated according to the NCI CTCAE v4.03. The incidence of DLT(s) during the first 6 weeks of treatment in Part 1 (the DLT evaluation period) and 4 weeks for Part 2 will be used.
Secondary Obj.:	1. Maximim Concentration (Cmax) 2. Time to Maximum Concentration (Tmax) 3. Trough Observed Plasma Concentration (Ctrough)

2.2 Background & Rationale

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

The purpose of this study is to evaluate the safety profile, tolerability, drug levels, drug effects, and preliminary efficacy of BMS-813160 alone or in combination with either chemotherapy or nivolumab or chemotherapy plus nivolumab in participants with metastatic colorectal and pancreatic cancers.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE1, PHASE2, Status: COMPLETED

Target N: 332

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Must have metastatic colorectal or pancreatic cancer
- * Eastern Cooperative Oncology Group (ECOG) performance status of ?1
- * Ability to swallow pills or capsules
- * Required to undergo mandatory pre and on-treatment biopsies
- * Adequate marrow function
- * Adequate other organ functions
- * Ability to comply with study visits, treatment, procedures, pharmacokinetic (PK) and pharmacodynamic (PD) sample collection, and required study follow-up

4.2 Exclusion Criteria

- * Histology other than adenocarcinoma (neuroendocrine or acinar cell)
- * Suspected, known, or central nervous system (CNS) metastases (Imaging required only if participants are symptomatic)
- * Active, known or suspected autoimmune disease
- * Condition requiring systemic treatment with either corticosteroids (> 10 mg daily prednisone equivalents) or other immunosuppressive medications within 14 days of study treatment administration
- * Interstitial lung disease that is symptomatic or may interfere with the detection or management of suspected treatment-related pulmonary toxicity
- * Prior treatment with cysteine-cysteine chemokine receptor 2 (CCR2) and/or cysteine-cysteine chemokine receptor 5 (CCR5) inhibitors, programmed death-1 receptor (PD-1), programmed death-ligand 1 \[PD(L)-1\] or cytotoxic T-lymphocyte antigen-4 (CTLA-4) antibodies
- * History of allergy to study treatments or any of its components of the study arm that participant is enrolling

Other protocol-defined inclusion/exclusion criteria apply

11. Study Identifiers & Governance

Primary ID: 22720114948792005

Registry Number: NCT03184870

Sponsor: Bristol Myers Squibb

Study Title: A Study of BMS-813160 in Combination With Chemotherapy or Nivolumab in

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Participants With Advanced Solid Tumors

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____