

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase Ib/II, Open-Label, Multicenter, Randomized Umbrella Study Evaluating the Efficacy and Safety of Multiple Immunotherapy-Based Treatment Combinations in Patients With Hormone Receptor-Positive HER2-Negative Breast Cancer (MORPHEUS-HR+ Breast Cancer)
Short Title:	A Study of Multiple Immunotherapy-Based Treatment Combinations in Hormone Receptor (HR)-Positive Human Epidermal Growth Factor Receptor 2 (HER2)-Negative Breast Cancer
Protocol Number:	347634327880628897
NCT Number:	NCT03280563
Sponsor Name:	Roche
Investigational Product:	tamoxifen
Phase:	PHASE1, PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Stage 1: Percentage of Participants With Objective Response: Objective response rate (ORR) was defined as the percentage of participants with an objective response of complete response (CR) or partial response (PR) as determined by the investigator according to Response Evaluation Criteria in Solid Tumors, Version 1.1 (RECIST v1.1). CR was defined as the disappearance of all target lesions and any pathological lymph nodes (whether target or non-target) that have a reduction in short axis to $\leq$ 10 millimeters (mm). PR was defined as at least a 30% decrease in the sum of diameters (SOD) of all target lesions, taking as reference the baseline SOD, in the absence of CR. Participants with missing or no response assessments were classified as non-responders. Percentages have been rounded off to the nearest whole number.
Secondary Obj.:	1. Stage 1: Progression-free Survival (PFS) 2. Stage 1: Clinical Benefit Rate (CBR) 3. Stage 1: Overall Survival (OS)

2.2 Background & Rationale

This study is designed to evaluate the efficacy, safety, and pharmacokinetics of several immunotherapy-based combination treatments in participants with inoperable locally advanced or metastatic HR-positive, HER2-negative breast cancer who have progressed during or following treatment with a cyclin-dependent kinase (CDK) 4/6 inhibitor in the first- or second-line setting, such as palbociclib, ribociclib, or abemaciclib. The study will be performed in two stages. During Stage 1, participants will be randomized to fulvestrant (control) or an atezolizumab-containing doublet or triplet combination. Those who experience disease progression, loss of clinical benefit, or unacceptable toxicity may be eligible to receive a new triplet combination treatment in Stage 2 until loss of clinical benefit or unacceptable toxicity. New

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

treatment arms may be added and/or existing treatment arms may be closed during the course of the study on the basis of ongoing clinical efficacy and safety as well as the current treatments available.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE1, PHASE2, Status: COMPLETED

Target N: 144

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

Not Provided

4.2 Exclusion Criteria

Not Provided

11. Study Identifiers & Governance

Primary ID: 347634327880628897

Registry Number: NCT03280563

Sponsor: Roche

Study Title: A Study of Multiple Immunotherapy-Based Treatment Combinations in Hormone Receptor (HR)-Positive Human Epidermal Growth Factor Receptor 2 (HER2)-Negative Breast Cancer

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____