

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 1b/2 Open-label Study Evaluating Different MK-6070 and Ifinatamab Deruxtecan (MK-2400)-Based Regimens in First-line Extensive Stage Small Cell Lung Cancer
Short Title:	A Clinical Study of Gocatamig (MK-6070) and Infinatamab Deruxtecan (MK-2400) in People With Small Cell Lung Cancer (MK-6070-003)
Protocol Number:	369951154487414396
NCT Number:	NCT07227597
Sponsor Name:	Merck
Investigational Product:	etoposide
Phase:	PHASE1, PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Number of Participants Who Experience an Adverse Event (AE): An AE is defined as any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. The number of participants who experience an AE will be reported. 2. Number of Participants Who Experience One or More Dose-Limiting Toxicities (DLTs): DLT will be defined as any drug-related AE observed during the DLT evaluation period (up to 21 days) that meets the protocol-specified DLT criteria. The number of participants who experience at least one DLT will be presented. 3. Number of Participants Who Discontinue Study Intervention Due to an AE: An AE is defined as any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. The number of participants who discontinue study intervention due to an AE will be reported.
Secondary Obj.:	1. Disease Control Rate (DCR) 2. Duration of Response (DOR) 3. Progression-Free Survival (PFS)

2.2 Background & Rationale

Researchers are looking for new ways to treat extensive-stage small cell lung cancer (ES-SCLC). ES-SCLC is a type of lung cancer that has spread throughout the lung, to the other lung, or to other parts of the body.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL,	Phase: PHASE1, PHASE2,	Status: NOT_YET_RECRUITING
-------------------	-----------------------	------------------------	----------------------------

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Target N: 170

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

The main inclusion criteria include but are not limited to the following:

- * Has a histologically or cytologically confirmed diagnosis of extensive-stage small cell lung cancer (ES-SCLC)
- * For participants receiving gocatamig + ifinatamab deruxtecan (I-DXd) in maintenance only:
 - * Completed 3 to 4 cycles of platinum + etoposide chemotherapy with concurrent approved anti-programmed cell death 1/Ligand 1 (anti PD-1/L1) as first line (1L) treatment of ES-SCLC within 6 weeks prior to enrollment
 - * No radiological disease progression per Response Evaluation Criteria In Solid Tumors version 1.1 (RECIST 1.1)
 - * No other prior systemic ES-SCLC therapy allowed
 - * Rechallenge therapy counts as an additional line and leads to exclusion
- * For participants receiving gocatamig + I-DXd in induction and maintenance, or gocatamig + I-DXd in induction followed by gocatamig + atezolizumab in maintenance, or carboplatin + etoposide + atezolizumab in induction followed by atezolizumab in maintenance: No prior systemic ES-SCLC treatment allowed
- * Applicable to all participants: prior limited-stage small cell lung cancer (SCLC) is allowed if > 6 months have passed since the end of previous therapy and progression
- * Must be able to provide a pretreatment archival tumor tissue sample or newly obtained core, incisional, or excisional biopsy of a tumor lesion not previously irradiated
- * Measurable disease by RECIST 1.1 as assessed by the local site investigator/radiology. Lesions situated in a previously irradiated area are considered measurable if growth has been shown in such lesions since the completion of radiation

4.2 Exclusion Criteria

The main exclusion criteria include but are not limited to the following:

- * Has pleural effusion, pericardial effusion, or ascites requiring recurrent drainage procedures
- * Has history of (noninfectious) pneumonitis/interstitial lung disease (ILD) that required steroids or has current pneumonitis/ILD, and or suspected pneumonitis/ILD
- * Has clinically severe pulmonary compromise resulting from intercurrent pulmonary illnesses
- * Has history of clinically significant intracranial bleeding or spinal cord bleeding
- * Has active neurologic paraneoplastic syndrome
- * Has history of coronary/peripheral artery bypass graft and/or any coronary/peripheral angioplasty or clinically significant cardiovascular disease such as myocardial infarction, symptomatic congestive heart failure (CHF), and/or uncontrolled cardiac arrhythmia within 6 months before the first dose of study intervention
- * Has other uncontrolled or significant protocol specified cardiovascular disease
- * Has history of arterial thrombosis within 6 months before the first dose of study intervention
- * Has chronic liver disease

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Has history of allogeneic tissue/solid organ transplant
- * Has history of leptomeningeal disease
- * Is infected with human immunodeficiency virus (HIV) and has a history of Kaposi's sarcoma and/or Multicentric Castleman's Disease
- * Has received prior therapy with an anti-programmed cell death protein 1 (anti-PD-1), anti-programmed cell death ligand 1 (anti-PD-L1), or anti-programmed cell death ligand 2 (anti-PD-L2) agent or with an agent directed to another stimulatory or coinhibitory T-cell receptor
- * Has received prior radiotherapy within 2 weeks of start of study intervention, or has radiation-related toxicities, requiring corticosteroids
- * Has known additional malignancy that is progressing or has required active treatment within the past 3 years
- * Has active autoimmune disease that has required systemic treatment in the past 2 years except replacement therapy
- * Has known active central nervous system (CNS) metastases and/or carcinomatous meningitis
- * Has major surgery within 4 weeks or minor surgery within 2 weeks of allocation (or first dose), or is anticipated to require a major surgical procedure during the study

11. Study Identifiers & Governance

Primary ID: 369951154487414396

Registry Number: NCT07227597

Sponsor: Merck

Study Title: A Clinical Study of Gocatumig (MK-6070) and Infinatamab Deruxtecan (MK-2400) in People With Small Cell Lung Cancer (MK-6070-003)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____