

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 1/2 Open-Label, Umbrella Platform Design Study of MK-2870 With Pembrolizumab (MK-3475) and Chemotherapy in Participants With 1L Locally Advanced Unresectable/Metastatic Gastroesophageal Adenocarcinoma (Gastric Adenocarcinoma, Gastroesophageal Junction Adenocarcinoma, and Esophageal Adenocarcinoma): Substudy 06C
Short Title:	Substudy 06C: A Study of Sacituzumab Tirumotecan (MK-2870) With Pembrolizumab (MK-3475) and Chemotherapy in Participants With First-Line Locally Advanced Unresectable/Metastatic Gastroesophageal Adenocarcinoma (MK-3475-06C/KEYMAKER-U06)
Protocol Number:	525113344818553890
NCT Number:	NCT06469944
Sponsor Name:	Merck
Investigational Product:	capecitabine
Phase:	PHASE1, PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Safety Lead-in Phase: Number of Participants Who Experience One or More Dose-Limiting Toxicities (DLTs): DLTs are defined as any drug-related adverse event (AE) according to the National Cancer Institute Common Terminology for Adverse Events (NCI CTCAE) Version 5.0, observed during the DLT evaluation period that results in a change to a given dose or a delay in initiating the next cycle. The percentage of participants who experience at least one DLT will be reported. 2. Safety Lead-in Phase: Number of Participants Who Experienced an Adverse Event (AE): An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. An AE can therefore be any unfavorable and unintended sign (including an abnormal laboratory finding), symptom, or disease (new or exacerbated) temporally associated with the use of a study intervention. 3. Safety Lead-in Phase: Number of Participants Who Discontinued Study Intervention Due to an AE: An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. An AE can therefore be any unfavorable and unintended sign (including an abnormal laboratory finding), symptom, or disease (new or exacerbated) temporally associated with the use of a study intervention.
Secondary Obj.:	1. Progression-Free Survival (PFS) per RECIST 1.1 as Assessed by BICR 2. Duration

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

of Response (DOR) Per RECIST 1.1 as Assessed by BICR 3. Overall Survival (OS)

2.2 Background & Rationale

This is a phase 1/2, multicenter, open-label umbrella platform study that will evaluate the safety and tolerability of sacituzumab tirumotecan with pembrolizumab and fluoropyrimidine chemotherapy for the first-line (1L) treatment of participants with locally advanced unresectable or metastatic human epidermal growth factor receptor 2 (HER2)-negative gastric, gastroesophageal junction, or esophageal adenocarcinoma.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE1, PHASE2, Status: RECRUITING

Target N: 130

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

The main inclusion criteria include but are not limited to the following:

- * Has histologically and/or cytologically confirmed diagnosis of previously untreated locally advanced unresectable or metastatic 1L gastroesophageal adenocarcinoma
- * Has gastroesophageal adenocarcinoma that is known to be human epidermal growth factor receptor 2 (HER2)/neu-positive are excluded. HER2 status is not required if HER2/neu testing is not mandatory per local standard of care (SOC)
- * Is not expected to require tumor resection during the treatment course
- * Has not had prior systemic therapy administered in the recurrent or metastatic setting
- * Has provided an archival tumor tissue sample or most recently obtained core, or incisional, or excisional biopsy for a tumor lesion
- * Participants who have adverse events (AEs) due to previous anticancer therapies must have recovered to <Grade 1 or baseline. Participants with endocrine-related AEs who are adequately treated with hormone replacement therapy are eligible
- * Has adequate organ function
- * Has measurable disease per Response Evaluation Criteria in Solid Tumors Version 1.1 (RECIST 1.1) as determined by the local site investigator/radiology assessment and verified by blind independent review committee (BICR)
- * Has Eastern Cooperative Oncology Group (ECOG) performance status of 0 or 1 assessed within 3 days before allocation/randomization.
- * Has a life expectancy of at least 6 months
- * Who are Hepatitis B surface antigen (HBsAg) positive are eligible if they have received hepatitis B virus (HBV) antiviral therapy for at least 4 weeks, and have undetectable HBV viral load prior to randomization.
- * Who has history of hepatitis C virus (HCV) infection are eligible if HCV viral load is undetectable at screening.
- * Human immunodeficiency virus (HIV)-infected participants must have well-controlled HIV on antiretroviral therapy (ART)

4.2 Exclusion Criteria

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

The main exclusion criteria include but are not limited to the following:

- * Has squamous cell or undifferentiated gastroesophageal cancer.
- * Has had previous therapy for locally advanced unresectable or metastatic gastric/GEJ/esophageal adenocarcinoma
- * Has experienced weight loss $\geq 20\%$ over 3 months before the first dose of study intervention
- * Has a history of documented severe dry eye syndrome, severe Meibomian gland disease and/or blepharitis, or severe corneal disease that prevents/delays corneal healing
- * Has Grade ≥ 2 peripheral neuropathy
- * Has active inflammatory bowel disease requiring immunosuppressive medication or previous history of inflammatory bowel disease
- * Has uncontrolled, significant cardiovascular disease or cerebrovascular disease within 6 months preceding study intervention
- * Has accumulation of pleural, ascitic, or pericardial fluid requiring drainage or diuretic drugs within 2 weeks prior to enrollment
- * HIV-infected participants with a history of Kaposi's sarcoma and/or Multicentric Castleman's Disease
- * Has received prior treatment with a trophoblast antigen 2 (TROP2)-targeted antibody-drug conjugate (ADC)
- * Has received prior treatment with a topoisomerase I inhibitor-based ADC and/or a topoisomerase I inhibitor-based chemotherapy
- * Has received prior systemic anticancer therapy including investigational agents within 4 weeks before the first dose of study intervention
- * Has received prior therapy with an anti-Programmed Cell Death Protein 1 (PD-1), anti-Programmed Cell Death-Ligand 1 (PD-L1), anti-Programmed Cell Death-Ligand 2 (PD-L2) agent or with an agent directed to another stimulatory or co-inhibitory T-cell receptor (TCR)
- * Has received prior radiotherapy within 2 weeks of start of study intervention, or has radiation related toxicities, requiring corticosteroids
- * Has received a live or live-attenuated vaccine within 30 days before the first dose of study intervention
- * Has received a strong inducer/inhibitor of CYP3A4 that cannot be discontinued
- * Has received an investigational agent or has used an investigational device within 4 weeks prior to study intervention administration
- * Has diagnosis of immunodeficiency or is receiving chronic systemic steroid therapy or any other form of immunosuppressive therapy within 7 days prior to the first dose of study intervention
- * Has known additional malignancy that is progressing or has required active treatment within the past 3 years
- * Has known active central nervous system (CNS) metastases and/or carcinomatous meningitis
- * Has Severe hypersensitivity (?Grade 3) to pembrolizumab, sacituzumab tirumotecan, or other biologic therapy, chemotherapy (ie, oxaliplatin, fluorouracil, capecitabine), leucovorin, levoleucovorin, or any of their excipients
- * Has active autoimmune disease that has required systemic treatment in the past 2 years
- * Has history of (noninfectious) pneumonitis/interstitial lung disease that required steroids or has current pneumonitis/interstitial lung disease
- * Has an active infection requiring systemic therapy
- * Has concurrent active hepatitis B (defined as Hepatitis B surface antigen [HBsAg] positive and/or detectable HBV

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

DNA) and hepatitis C virus (defined as anti-hepatitis C virus [HCV] Ab positive and detectable HCV ribonucleic acid [RNA] infection

- * Has GI obstruction, poor oral intake, or difficulty in taking oral medication
- * Has poorly controlled diarrhea
- * Has had a major surgery or significant traumatic injury within 4 weeks before the first dose of study intervention
- * Has history of allogeneic tissue/solid organ transplant
- * Have not adequately recovered from major surgery or have ongoing surgical complications

11. Study Identifiers & Governance

Primary ID: 525113344818553890

Registry Number: NCT06469944

Sponsor: Merck

Study Title: Substudy 06C: A Study of Sacituzumab Tirumotecan (MK-2870) With Pembrolizumab (MK-3475) and Chemotherapy in Participants With First-Line Locally Advanced Unresectable/Metastatic Gastroesophageal Adenocarcinoma (MK-3475-06C/KEYMAKER-U06)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____