

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 1/2 Open-Label, Umbrella Platform Design Study to Evaluate the Safety and Efficacy of MK-2870 Plus Paclitaxel as the Second-Line Treatment of Participants With Advanced/Metastatic Gastroesophageal Adenocarcinoma: Substudy 06D
Short Title:	Substudy 06D: Combination Therapies in Second Line (2L) Gastroesophageal Adenocarcinoma (MK-3475-06D/Keymaker-U06)
Protocol Number:	737280407558717956
NCT Number:	NCT06445972
Sponsor Name:	Merck
Investigational Product:	paclitaxel
Phase:	PHASE1, PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Percentage of Participants who Experience Dose Limiting Toxicities (DLTs) During the Safety Lead-In Phase: DLTs are defined as any drug-related adverse event (AE) according to the National Cancer Institute Common Terminology for Adverse Events (NCI CTCAE) Version 5.0, observed during the DLT evaluation period that results in a change to a given dose or a delay in initiating the next cycle. The percentage of participants who experience at least one DLT will be presented. 2. Percentage of Participants who Experience an Adverse Event (AE) During the Safety Lead-In Phase: An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. The percentage of participants who discontinued study intervention due to an AE will be presented. 3. Percentage of Participants who Discontinue Study Intervention Due to an AE During the Safety Lead-In Phase: An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. The percentage of participants who discontinued study intervention due to an AE will be presented.
Secondary Obj.:	1. Progression Free Survival (PFS) 2. Duration of Response (DOR) 3. Overall Survival (OS)

2.2 Background & Rationale

This is a phase 1/2 multicenter, open-label umbrella platform study that will evaluate the safety and efficacy of sacituzumab tirumotecan (MK-2870) plus paclitaxel versus ramucirumab plus paclitaxel, for the treatment of participants with advanced or metastatic gastric adenocarcinoma, gastroesophageal junction (GEJ) adenocarcinoma, or esophageal

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

adenocarcinoma who have failed 1 prior line of therapy. This is an estimation study, and no formal hypothesis testing will be performed.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE1, PHASE2, Status: RECRUITING

Target N: 90

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

The main inclusion criteria include but are not limited to the following:

- * Has histologically and/or cytologically confirmed diagnosis of previously treated, 2L (received first line (1L) treatment) gastric adenocarcinoma, GEJ adenocarcinoma, or esophageal adenocarcinoma
- * Has metastatic disease or locally advanced, unresectable disease
- * Has experienced documented objective radiographic or clinical disease progression during or after 1L therapy containing any platinum/fluoropyrimidine doublet with or without immunotherapy
- * Participants with gastroesophageal adenocarcinoma that is known to be human epidermal growth factor receptor 2 (HER2)/neu positive are excluded. Participants with unknown HER2 status are eligible.
- * Has provided an archival tumor tissue sample or most recently obtained core, incisional, or excisional biopsy of a tumor lesion
- * AEs due to previous anticancer therapies must be ?Grade 1 or baseline (except alopecia and vitiligo). Endocrine-related AEs adequately treated with hormone replacement are acceptable.
- * Has Eastern Cooperative Oncology Group performance status of 0 or 1
- * Has a life expectancy of at least 3 months
- * Participants who are hepatitis B surface antigen (HBsAg) positive are eligible if they have received Hepatitis B Virus (HBV) antiviral therapy for at least 4 weeks, and have undetectable HBV viral load prior to randomization.
- * Participants with history of Hepatitis C Virus (HCV) infection are eligible if HCV viral load is undetectable at screening.
- * Human Immunodeficiency Virus (HIV)-infected participants must have well controlled HIV on antiretroviral therapy.

4.2 Exclusion Criteria

The main exclusion criteria include but are not limited to the following:

- * Has squamous cell or undifferentiated gastroesophageal cancer
- * Has experienced weight loss \>20% over 3 months before the first dose of study intervention
- * Has history of documented severe dry eye syndrome, severe Meibomian gland disease and/or blepharitis, or severe corneal disease that prevents/delays corneal healing
- * Has Grade ?2 peripheral neuropathy
- * Has active inflammatory bowel disease requiring immunosuppressive medication or previous history of inflammatory bowel disease
- * Has a serious or nonhealing wound or peptic ulcer or bone fracture within 28 days prior to randomization

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Has a bowel obstruction, history or presence of inflammatory enteropathy or extensive intestinal resection (hemicolectomy or extensive small intestine resection with chronic diarrhea)
- * Has uncontrolled, significant cardiovascular disease or cerebrovascular disease
- * Has experienced any arterial thrombotic event, including myocardial infarction, unstable angina, cerebrovascular accident, or transient ischemic attack, within 6 months prior to randomization
- * Has uncontrolled arterial hypertension $\geq 150/\geq 90$ mm mercury (Hg)
- * Has accumulation of pleural, ascitic, or pericardial fluid requiring drainage or diuretic drugs within 2 weeks prior to enrollment
- * Has undergone major surgery within 28 days prior to randomization, or central venous access device placement within 7 days prior to randomization or planned major surgery following initiation of study treatment
- * Is receiving therapeutic anticoagulation with warfarin, low-molecular weight heparin or similar agents
- * Is receiving chronic therapy with nonsteroidal anti-inflammatory agents (NSAIDs) or other antiplatelet agents
- * Has a history of deep vein thrombosis, pulmonary embolism, or any other significant thromboembolism (venous port or catheter thrombosis or superficial venous thrombosis are not considered "significant") during the 3 months prior to randomization
- * Has significant bleeding disorders, vasculitis, or had a significant bleeding episode from the gastrointestinal (GI) tract within 3 months prior to study entry
- * Has history of GI perforation and/or fistulae within 6 months prior to randomization
- * HIV-infected participants with a history of Kaposi's sarcoma and/or Multicentric Castleman's Disease
- * Has received prior treatment with a trophoblast cell-surface antigen 2 (TROP2)-targeted antibody-drug conjugate (ADC), topoisomerase 1 inhibitor-based ADC and/or a topoisomerase 1 inhibitor-based chemotherapy
- * Has received any previous systemic therapy targeting vascular endothelial growth factor (VEGF) or the vascular endothelial growth factor receptor (VEGFR) signaling pathways
- * Has received prior systemic anticancer therapy including investigational agents within 4 weeks before the first dose of study drug intervention
- * Has received prior radiotherapy within 2 weeks of start of study intervention, or has radiation-related toxicities, requiring corticosteroids
- * Has received a live or live-attenuated vaccine within 30 days before the first dose of study intervention. Administration of killed vaccines is allowed
- * Has received an investigational agent or has used an investigational device within 4 weeks prior to study intervention administration
- * Has a known additional malignancy that is progressing or has required active treatment within the past 3 years. Basal cell carcinoma of the skin, squamous cell carcinoma of the skin, or carcinoma in situ, excluding carcinoma in situ of the bladder, that have undergone potentially curative therapy are not excluded.
- * Has known active central nervous system (CNS) metastases and/or carcinomatous meningitis
- * Has an active infection requiring systemic therapy
- * Has a concurrent active HBV (defined as HBsAg positive and/or detectable HBV deoxyribonucleic acid) and HCV (defined as anti-HCV antibody positive and detectable HCV ribonucleic acid) infection
- * History of (noninfectious) pneumonitis/interstitial lung disease that required steroids or has current pneumonitis/interstitial lung disease

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

* Has severe hypersensitivity (Grade ?3) to sacituzumab tirumotecan, any of its excipients and/or to another biologic therapy

* Has not adequately recovered from major surgery or have ongoing surgical complications

11. Study Identifiers & Governance

Primary ID: 737280407558717956

Registry Number: NCT06445972

Sponsor: Merck

Study Title: Substudy 06D: Combination Therapies in Second Line (2L) Gastroesophageal Adenocarcinoma (MK-3475-06D/Keymaker-U06)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____