

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase Ib/II, Multicenter Study of the Combination of LEE011 and BYL719 With Letrozole in Adult Patients With Advanced ER+ Breast Cancer
Short Title:	Study of LEE011, BYL719 and Letrozole in Advanced ER+ Breast Cancer
Protocol Number:	782986010198506109
NCT Number:	NCT01872260
Sponsor Name:	Novartis
Investigational Product:	letrozole
Phase:	PHASE1, PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Incidence of Dose limiting toxicities (DLTs) - Phase Ib only 2. Safety and tolerability: Adverse Events (AEs), serious AEs (SAEs), changes in hematology and chemistry values, vital signs, electrocardiograms (ECGs), dose interruptions, reductions and dose intensity. 3. PK profiles of LEE011 and letrozole: To characterize PK profiles of LEE011 and Letrozole.
Secondary Obj.:	1. Safety and tolerability of LEE011 in combination with letrozole, BYL719 in combination with letrozole, and the triple combination of LEE011 +BYL719 with letrozole 2. Plasma concentration-time profiles of LEE011, BYL719 and letrozole 3. Overall Response Rate (ORR)

2.2 Background & Rationale

The purpose of this trial is to inform the future clinical development of the two investigational agents in ER+ breast cancer, LEE011 (CDK4/6 inhibitor) and BYL719 (PI3K-alpha inhibitor).

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL,	Phase: PHASE1, PHASE2,	Status: ACTIVE_NOT_RECRUITING
Target N:	255		

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Postmenopausal, Estrogen-receptor positive and/or Progesterone-receptor positive breast cancer
- * Phase Ib dose escalation only: Any number of prior lines of endocrine therapy is allowed with the exception of cytotoxic therapy which is limited to one prior line administered in the advanced (metastatic or locally advanced) setting.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Phase Ib dose expansions Arms 1, 2 and 3
- * No prior systemic treatment in the advanced (metastatic or locally advanced) setting with the exception of treatment with letrozole for a maximum of one month prior to starting study treatment.
- * Patients who received (neo)adjuvant therapy for breast cancer are eligible. Prior therapy with letrozole or anastrozole in the (neo)adjuvant setting is permitted if the disease-free interval is greater than 12 months from the completion of treatment.

4.2 Exclusion Criteria

- * HER2-overexpression in the patient's tumor tissue
- * Patients with active CNS or other brain metastases
- * Major surgery within 2 weeks
- * Acute or chronic pancreatitis
- * Bilateral diffuse lymphangitic carcinomatosis
- * Another malignancy within 3 years
- * Receiving hormone replacement therapy that cannot be discontinued
- * Impaired cardiac function
- * Patients with clinically manifest diabetes mellitus (treated and/or clinical signs or with fasting glucose ≥ 126 mg/dL / 7.0 mmol/L or hemoglobin A1c $\geq 6.5\%$), history of gestational diabetes mellitus or documented steroid-induced diabetes mellitus.
- * Other protocol-defined inclusion/exclusion criteria may apply

11. Study Identifiers & Governance

Primary ID: 782986010198506109
Registry Number: NCT01872260
Sponsor: Novartis
Study Title: Study of LEE011, BYL719 and Letrozole in Advanced ER+ Breast Cancer

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____