

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 1b/2 Open-label, Multicenter Study to Evaluate the Safety and Efficacy of Raludotatug Deruxtecan With or Without Other Anticancer Investigational Agents in Participants With High-grade Serous Epithelial Ovarian, Primary Peritoneal, or Fallopian Tube Cancer Who Have Relapsed After Prior Platinum-based Chemotherapy
Short Title:	A Clinical Study of Raludotatug Deruxtecan in People With Ovarian Cancer (MK-5909-003)
Protocol Number:	803265315070577003
NCT Number:	NCT06843447
Sponsor Name:	Merck
Investigational Product:	carboplatin
Phase:	PHASE1, PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Part 1: Number of Participants Who Experience a Dose-limiting Toxicity (DLT) Per Common Terminology Criteria for Adverse Events, Version 5.0 (CTCAE v5.0): DLTs are defined as toxicities during the DLT evaluation period that are assessed by the investigator to be possibly, probably, or definitely related to study treatment and include: Grade 4 thrombocytopenia of any duration or Grade 3 thrombocytopenia lasting ≥7 days; Grade 3 or higher thrombocytopenia associated with clinically significant bleeding; Grade 4 lymphocytopenia lasting ≥14 days; Grade 4 anemia of any duration; any other Grade 4 hematologic toxicity lasting ≥7 days; febrile neutropenia Grade 3 or Grade 4 meeting pre-specifications; pre-specified hepatic organ toxicities; all Grade 3 or higher other nonhematologic toxicities except those pre-specified; other pre-specified nonhematologic toxicities; any delay in treatment with the planned dose of ≥21 days or discontinuation of treatment due to a toxicity during the DLT evaluation period, or Grade 5 toxicity. The number of participants with DLTs will be reported. 2. Part 1: Number of Participants with One or More Adverse Events (AEs): An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. The number of participants with one or more AEs will be reported. 3. Part 1: Number of Participants who Discontinue Study Intervention Due to an AE: An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. The number of participants who discontinue study intervention due to an AE will be reported.
Secondary Obj.:	1. Part 1: Objective Response Rate (ORR) 2. Part 2: Duration of Response (DOR) 3.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Part 2: Progression-free Survival (PFS)

2.2 Background & Rationale

Researchers are looking for other ways to treat relapsed high-grade serous ovarian cancer. Relapsed means the cancer came back after treatment. High-grade means the cancer cells grow and spread quickly. Serous means the cancer started in the cells that cover the ovaries, the lining of the belly, or in the fallopian tubes.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE1, PHASE2, Status: RECRUITING

Target N: 280

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Has pathologically documented diagnosis of high-grade serous epithelial ovarian cancer, primary peritoneal cancer, or fallopian tube cancer
- * Has measurable disease per Response Evaluation Criteria In Solid Tumors 1.1
- * Participants in Cohort A-1 Arm 2 and Arm 3: Has relapsed disease after 1 to 3 prior lines of therapy and radiographic evidence of disease progression ≥6 months (≥180 days) after the last dose of platinum-based therapy (ie, platinum-sensitive disease).
- * Participants in Cohort B-1 and Cohort B-2: Has relapsed disease after 1 to 3 prior lines of therapy and radiographic evidence of disease progression <6 months (<180 days) after the last dose of platinum-based therapy (ie, platinum-resistant disease).
- * Participants in Cohort B-1 and Cohort B-2: Is a candidate for bevacizumab treatment
- * Has provided tumor tissue from a core or excisional biopsy of a tumor lesion not previously irradiated
- * Has an Eastern Cooperative Oncology Group performance status of 0 to 1 assessed within 7 days before allocation
- * Human immunodeficiency virus (HIV)-infected participants must have well controlled HIV on anti-retroviral therapy
- * Participants who are hepatitis B surface antigen positive are eligible if they have received hepatitis B virus (HBV) antiviral therapy for at least 4 weeks, and have undetectable HBV viral load prior to allocation
- * Participants with a history of hepatitis C virus (HCV) infection are eligible if HCV viral load is undetectable at screening
- * Participants in Cohort C-1 and Cohort D: Has relapsed disease after 1 prior line of therapy, radiographic evidence of disease progression ≥6 months (≥180 days) after the last dose of platinum-based therapy (ie, platinum-sensitive disease) and progressed during prior treatment with PARPi in the first-line setting

4.2 Exclusion Criteria

- * Has any of the following within 6 months before allocation: cerebrovascular accident, transient ischemic attack, or other arterial thromboembolic event
- * Has uncontrolled or significant cardiovascular disease
- * Has clinically severe pulmonary compromise resulting from intercurrent pulmonary illnesses including, but not limited to, any underlying pulmonary disorder, and any autoimmune, connective tissue, or inflammatory disorders with potential pulmonary involvement, or prior pneumonectomy
- * Has ≥Grade 2 peripheral neuropathy

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Has received prior treatment with cadherin-6-targeted agents
- * Has received prior systemic anticancer therapy including investigational agents within 4 weeks or 5 half-lives (whichever is shorter) before allocation
- * Has received prior radiotherapy within 2 weeks of the start of study intervention, or has radiation-related toxicities, requiring corticosteroids
- * Receives chronic steroid treatment
- * Has known additional malignancy that is progressing or has required active treatment within the past 3 years
- * Has known active CNS metastases and/or carcinomatous meningitis
- * Has history of (noninfectious) pneumonitis/interstitial lung disease (ILD) that required steroids or has current pneumonitis/ILD
- * Has active infection requiring systemic therapy
- * HIV-infected participants with a history of Kaposi's sarcoma and/or Multicentric Castleman's Disease

11. Study Identifiers & Governance

Primary ID: 803265315070577003

Registry Number: NCT06843447

Sponsor: Merck

Study Title: A Clinical Study of Raludotatug Deruxtecan in People With Ovarian Cancer (MK-5909-003)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____