

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	HERTHENA-Breast-01: A Phase 1b/2, Multicenter, Open-label, Dose-Finding Study to Evaluate the Safety and Antitumor Activity of Patritumab Deruxtecan in Participants With HER2 Positive Unresectable Locally Advanced Breast Cancer or Metastatic Breast Cancer
Short Title:	Study of Patritumab Deruxtecan With Other Anticancer Agents in Participants With HER2 Positive Breast Cancer That Has Spread and Cannot Be Surgically Removed (MK-1022-009)
Protocol Number:	805073040590599769
NCT Number:	NCT06686394
Sponsor Name:	Merck
Investigational Product:	trastuzumab
Phase:	PHASE1, PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Number of Participants Experiencing Dose-Limiting Toxicity (DLT): DLT will be defined as any drug-related AE observed during the DLT evaluation period that results in a change to a given dose or a delay in initiating the next cycle. The number of participants who experience a DLT will be presented. 2. Number of Participants with One or More Adverse Events (AEs): An AE is defined as any unfavorable and unintended sign, symptom, disease, or worsening of preexisting condition temporally associated with study treatment and irrespective of causality to study treatment. The number of participants who experience an AE will be presented. 3. Number of Participants who Discontinue Study Intervention Due to an AE: An AE is defined as any unfavorable and unintended sign, symptom, disease, or worsening of preexisting condition temporally associated with study treatment and irrespective of causality to study treatment. The number of participants who discontinue study treatment due to an AE will be presented.
Secondary Obj.:	1. Maximum Plasma Concentration (Cmax) of Patritumab Deruxtecan Antibody-Drug Conjugate (ADC) 2. Trough Concentration (Ctrough) of Patritumab Deruxtecan ADC 3. Area Under the Plasma Concentration-Time Curve (AUC) of Patritumab Deruxtecan ADC

2.2 Background & Rationale

Researchers want to learn if patritumab deruxtecan (MK-1022) can treat certain breast cancers. The breast cancers being studied are HER2 positive unresectable locally advanced or metastatic (the cancer has spread to other parts of

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

the body). The goals of this study are to learn:

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE1, PHASE2, Status: RECRUITING

Target N: 81

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

The main inclusion criteria include but are not limited to the following:

- * Has histologically confirmed HER2+ locally advanced unresectable breast cancer or metastatic breast cancer
- * Human immunodeficiency virus (HIV)-infected participants must have well-controlled HIV on antiretroviral therapy (ART)
- * Participants who are hepatitis B surface antigen (HBsAg) positive are eligible if they have received HBV antiviral therapy for at least 4 weeks, and have undetectable hepatitis B virus (HBV) viral load before allocation
- * Participants with history of hepatitis C virus (HCV) infection are eligible if HCV viral load is undetectable
- * Eastern Cooperative Oncology Group (ECOG) performance status 0 to 1 within 7 days before start of study intervention

Arm 1:

- * Has received at least a minimum of 2 and a maximum of 5 prior lines of anti-HER2 therapy in the locally advanced or metastatic setting
- * Had disease progression on or after any previous trastuzumab deruxtecan (T-DXd) treatment

Arm 2:

-Has received no more than 5 prior lines of anti-HER2 therapy in the locally advanced or metastatic setting

Arm 3:

-Has received and had disease progression from T-DXd treatment in any setting and a maximum of 3 prior lines of anti-HER2 therapy in the locally advanced or metastatic setting. T-DXd must be the most recent therapy received before enrollment.

4.2 Exclusion Criteria

The main exclusion criteria include but are not limited to the following:

- * Uncontrolled or significant cardiovascular disease
- * History of (noninfectious) pneumonitis/interstitial lung disease (ILD) that required steroids or has current

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

pneumonitis/interstitial lung disease

- * Has clinically severe respiratory compromise
- * Has any history of or evidence of any current leptomenigeal disease
- * Has clinically significant corneal disease
- * Evidence of ongoing uncontrolled systemic bacterial, fungal, or viral infection
- * HIV infected participants with a history of Kaposi's sarcoma and/or Multicentric Castleman's Disease
- * Known additional malignancy that is progressing or has required active treatment within the past 3 years
- * Evidence of spinal cord compression or brain metastases
- * Has an active infection requiring systemic therapy
- * Concurrent active HBV and HCV infection
- * Has had major surgical procedure (excluding placement of vascular access) less than 28 days

Arm 3 ONLY

↳ Has received prior treatment with tucatinib, lapatinib, or neratinib, or any investigational HER2-targeted tyrosine kinase inhibitors in the locally advanced or metastatic setting

11. Study Identifiers & Governance

Primary ID:	805073040590599769
Registry Number:	NCT06686394
Sponsor:	Merck
Study Title:	Study of Patritumab Deruxtecan With Other Anticancer Agents in Participants With HER2 Positive Breast Cancer That Has Spread and Cannot Be Surgically Removed (MK-1022-009)

15. Sign-off & Approval

Principal Investigator: _____	Date: _____
Clinical Operations Lead: _____	Date: _____
Lead Statistician: _____	Date: _____