

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Three-part, Phase I/II Study to Investigate the Safety, Tolerability, Pharmacokinetics, and Efficacy of Zifibancimig Following Intravitreal Administration of Multiple Ascending Doses and Continuous Delivery From the Port Delivery in Patients With Neovascular Age-related Macular Degeneration
Short Title:	Study of Zifibancimig in Participants With Neovascular Age-related Macular Degeneration
Protocol Number:	947170500794734699
NCT Number:	NCT04567303
Sponsor Name:	Roche
Investigational Product:	ranibizumab
Phase:	PHASE1, PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Percentage of Participants With Ocular and Systemic (Non-ocular) Adverse Events (AEs) 2. Percentage of Participants With Ocular AEs During the Post-operative and Follow-up Periods 3. Percentage of Participants With Adverse Events of Special Interest (AESIs) Including Ocular AESIs
Secondary Obj.:	1. Maximum Observed Concentration (Cmax) of Zifibancimig in Blood and Aqueous Humor (AH) 2. Time of Maximum Concentration Observed (Tmax) of Zifibancimig in Blood and AH 3. Concentration at the End of a Dosing Interval Before the Next Dose Administration (Ctrough) of Zifibancimig in Blood and AH

2.2 Background & Rationale

This is a first in-human study to investigate the safety, tolerability and efficacy of zifibancimig administered through intravitreal (IVT) injections and via the Port Delivery (PD) implant in participants with neovascular age-related macular degeneration (nAMD).

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE1, PHASE2, Status: RECRUITING
Target N:	251

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

* Willing to allow AH collection

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Part 1 and Part 2 Ocular Inclusion Criteria for Study Eye:

- * Choroidal neovascularization (CNV) exclusively due to age-related macular degeneration (AMD)
- * Anti-vascular endothelial growth factor (VEGF) or anti-VEGF/Angiopoietin-2 (Ang-2) IVT treatment-naïve, or pre-treated with anti-VEGF or anti-VEGF/Ang-2 no less than two months prior to Day 1
- * Sufficiently clear ocular media and adequate pupillary dilatation to allow for analysis and grading by the central reading center of fundus photography (FP), fluorescein angiography (FA), fundus autofluorescence (FAF), and spectral domain optical coherence tomography (SD-OCT) images
- * Decreased BCVA attributable primarily to nAMD, with BCVA letter score of 78 to 34 letters (inclusive) on ETDRS-like charts at screening. In case both eyes of a participant are eligible, the eye with the lower BCVA score should become the study eye

Part 3 Ocular Inclusion Criteria for Study Eye:

- * CNV exclusively due to AMD
- * Diagnosis of nAMD within 36-months prior to the screening visit
- * Previous treatment with at least one IVT anti-VEGF or anti-VEGF/Ang-2 administrations IVT for nAMD. The last IVT administration must have occurred at least 21 days prior to the screening visit
- * Demonstrated response to prior IVT anti-VEGF or anti-VEGF/Ang-2 treatment since diagnosis
- * Availability of historical VA data prior to the first anti-VEGF or anti-VEGF/Ang-2 treatment for nAMD
- * Sufficiently clear ocular media and adequate pupillary dilatation to allow for analysis and grading
- * Decreased BCVA attributable primarily to nAMD with letter score of 78 to 34 letters (inclusive) or better on ETDRS-like charts

Ocular Exclusion Criteria for Study Eye:

- * History of vitrectomy surgery, submacular surgery, other intraocular surgery, or any planned surgical intervention during the study period
- * Cataract surgery without complications within three months preceding the screening visit or planned during the study period
- * Aphakia or absence of the posterior capsule. Previous violation of the posterior capsule is also an exclusion criterion, unless it occurred as a result of yttrium-aluminum garnet laser posterior capsulotomy in association with prior, posterior chamber intraocular lens implantation
- * Prior macular treatment with verteporfin, external beam radiation therapy, transpupillary thermotherapy, or any type of laser photocoagulation
- * Prior treatment with IVT corticosteroids or implant (e.g., triamcinolone, ozurdex, iluvien)
- * Subretinal hemorrhage >50% of the total lesion area and/or involving the fovea
- * Subfoveal fibrosis or subfoveal atrophy
- * Retinal pigment epithelial tear involving the macula

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * History of vitreous hemorrhage, rhegmatogenous retinal detachment, glaucoma-filtering surgery, tube shunts, or microinvasive glaucoma surgery, and corneal transplant
- * History of rhegmatogenous retinal tears or peripheral retinal breaks within three months prior to the screening visit
- * Actual or history of myopia ≥ -8 diopters
- * Uncontrolled ocular hypertension or glaucoma (defined as intraocular pressure (IOP) ≥ 25 millimeters of mercury (mm Hg) or a cup to disc ratio ≥ 0.8 , despite treatment with antiglaucoma medication) and any such condition the Investigator determines may require a glaucoma-filtering surgery during a participant's participation in the study
- * Concurrent intraocular conditions (e.g., cataract, diabetic retinopathy, epiretinal membrane with traction, macular hole) that, in the opinion of the Investigator, could either: require medical or surgical intervention during the study period to prevent or treat visual loss that might result from that condition; or likely contribute to loss of BCVA over the study period if allowed to progress untreated; or preclude any visual improvement due to substantial structural damage
- * Concurrent conjunctival, Tenon's capsule, and/or scleral condition in the supero-temporal quadrant of the eye (e.g., scarring, thinning, mass) that may affect the implantation, subsequent tissue coverage, and refill-exchange procedure of the PD implant
- * Prior treatment with any medication for geographic atrophy during the last 3 months prior to screening
- * Prior treatment with any anti-VEGF-C or anti-VEGF-D inhibitors

Exclusion Criteria for Fellow Eye

- * BCVA letter score using ETDRS charts of ≤ 34 letters
- * Treatment with IVT anti-VEGF or anti-VEGF/Ang-2 agents within one week prior to Day 1 (concurrent treatment with SUSVIMOTM in the fellow eye is not exclusionary)

Exclusion Criteria for Either Eye

- * CNV due to causes other than nAMD, such as ocular histoplasmosis, trauma, pathological myopia, angioid streaks, choroidal rupture, uveitis or central serous chorioretinopathy
- * Prior participation in a clinical trial involving anti-VEGF drugs within six months prior to the screening visit, other than ranibizumab, aflibercept, or faricimab including approved biosimilars
- * Active intraocular inflammation (grade trace or above), infectious conjunctivitis, keratitis, scleritis, or endophthalmitis
- * History of uveitis, including history of any intraocular inflammation following intravitreal therapy
- * Prior treatment with brodalumab
- * Prior gene therapy for nAMD

4.2 Exclusion Criteria

Not Provided

11. Study Identifiers & Governance

Primary ID:	947170500794734699
Registry Number:	NCT04567303
Sponsor:	Roche

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Study Title: Study of Zifibancimig in Participants With Neovascular Age-related Macular Degeneration

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____