

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase Ib/II Open Label Dose Confirmation, Proof of Concept Study of Siremadlin in Combination With Venetoclax Plus Azacitidine in Unfit Adult AML Participants Who Responded Sub-optimally to First-line Venetoclax Plus Azacitidine Treatment and in Participants With Newly Diagnosed Unfit AML Presenting With High-risk Clinical Features
Short Title:	A Study of Siremadlin in Combination With Venetoclax Plus Azacitidine in Adult Participants With Acute Myeloid Leukemia (AML) Who Are Ineligible for Chemotherapy.
Protocol Number:	1027377931985192537
NCT Number:	NCT05155709
Sponsor Name:	Novartis
Investigational Product:	azacitidine
Phase:	PHASE1
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:

1. Percentage of participants with Dose Limiting Toxicities (DLTs) as per investigator assessment reported during the first cycle (separately in Arm 1 & Arm 2): The objective for the safety run-in is to determine the recommended dose of siremadlin in combination with venetoclax plus azacitidine to be explored further in the expansion part separately in Arm 1 and Arm 2. 2. Percentage of participants treated at the recommended dose for expansion, achieving a complete remission (CR) as per investigator assessment (Arm 1 only): The objective is to evaluate preliminary efficacy of siremadlin at the determined recommended dose for expansion (RDE) when administered in combination with venetoclax and azacitidine.

This endpoint will not be analyzed since the recommended dose was not determined due to early termination.

Secondary Obj.:

1. Percentage of participants treated at the recommended dose for expansion, achieving CR as per investigator assessment (Arm 2 only; for Arm 1 assessment of CR is a primary outcome measure)) 2. Time of the date of the first documented CR to the date of the first documented relapse or death due to any cause, whichever occurs first (Arm 1 and Arm 2 separately) 3. Percentage of participants achieving CR or complete remission with partial hematological recovery (CRh) and percentage of participants achieving CR or complete remission with incomplete hematological recovery (CRi) (Arm 1 and Arm 2)

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

2.2 Background & Rationale

A study of siremadlin in combination with venetoclax plus azacitidine in adult participants with AML who are ineligible for chemotherapy.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE1, Status: TERMINATED

Target N: 14

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

↳ Age at the date of signing the informed consent form (ICF): Arm 1 and Arm 2: ? 18 years

↳ Participants diagnosed with AML based on WHO 2016 classification (Arber et al 2016) who are ineligible for standard induction chemotherapy and: Arm 1 : have received at least 2 cycles and not more than 4 cycles of first-line venetoclax plus azacitidine treatment and have not achieved a CR, CRi, CRh or MLFS.

Arm 2 : newly diagnosed AML with adverse genetic risk stratification (according to ELN 2022) (except TP53 mutation positive participants).

* Participant (in both arms) must be considered ineligible for standard of care intensive induction chemotherapy defined by the following:

* 75 years of age; OR

* 18 to 74 years of age with at least one of the following co-morbidities: Eastern Cooperative Oncology Group (ECOG) Performance Status of 2 or 3; Cardiac history of congestive heart failure (CHF) requiring treatment or Ejection Fraction ? 50% or chronic stable angina; DLCO ? 65% or FEV1 ? 65%.

* Participants must have an ECOG performance status:

0 to 2 for participants ? 75 years of age. OR 0 to 3 for participants ? 18 to 74 years of age.

* WBC \< 25x10⁹/L

* AST and ALT ? 3 × ULN

* Estimated Glomerular Filtration Rate (eGFR)? 60 mL/min/1.73 m²

4.2 Exclusion Criteria

* Prior exposure to MDM2-inhibitor therapy at any time.

* Participants with TP53 mutation positive.

* Participants with del17p.

* Participants with AML-M3 / APL (Acute promyelocytic leukemia) with PML-RARA (Promyelocytic leukemia/retinoic acid receptor alpha) or with AML secondary to Down's syndrome.

* Participants treated with FLT3 inhibitors for AML indication are not eligible.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Participants who require treatment with moderate or strong CYP3A4 inducers within 14 days prior to starting study treatment, or are expected to receive moderate or strong CYP3A4 inducers during the entire study
- * Participants who require treatment with substrates of CYP3A4/5 with a narrow therapeutic index.

Other protocol-defined inclusion/exclusion criteria may apply at the end

11. Study Identifiers & Governance

Primary ID: 1027377931985192537

Registry Number: NCT05155709

Sponsor: Novartis

Study Title: A Study of Siremadlin in Combination With Venetoclax Plus Azacitidine in Adult Participants With Acute Myeloid Leukemia (AML) Who Are Ineligible for Chemotherapy.

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____