

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Randomized, Double-Blind, Placebo-Controlled Clinical Trial to Assess the Safety, Tolerability, and Pharmacokinetics of MK-1167 Administered to Patients With Alzheimer's Disease Receiving Stable Donepezil Treatment
Short Title:	Efficacy and Safety of MK-1167 in Participants With Alzheimer's Disease Dementia Taking Stable Donepezil Treatment (MK-1167-007)
Protocol Number:	1089974905016072149
NCT Number:	NCT06285240
Sponsor Name:	Merck
Investigational Product:	donepezil
Phase:	PHASE1
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Number of Participants Who Experienced an Adverse Event (AE): An AE was any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. An AE can therefore be any unfavorable and unintended sign (including an abnormal laboratory finding), symptom, or disease (new or exacerbated) temporally associated with the use of a study intervention. The number of participants who experienced an AE were reported. 2. Number of Participants Who Discontinued Study Treatment Due to an AE: An AE was any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. An AE can therefore be any unfavorable and unintended sign (including an abnormal laboratory finding), symptom, or disease (new or exacerbated) temporally associated with the use of a study intervention. The number of participants who discontinued study treatment due to an AE were reported.
Secondary Obj.:	1. Panel A: Area Under the Plasma Concentration-Time Curve From 0 to 24 Hours (AUC0-24) After Administration of 6mg of MK-1167 2. Panel A: AUC0-24 After Administration of 3mg of MK-1167 3. Panel B: AUC0-24 After Administration of 6mg of MK-1167

2.2 Background & Rationale

The main purpose of this study is to assess the safety and efficacy of MK-1167 administered to participants with Alzheimer's Disease (AD) receiving stable Donepezil treatment.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE1, Status: COMPLETED

Target N: 28

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Reports a history of cognitive and functional decline with gradual onset and slow progression for at least 1 year before Screening, that is either corroborated by an informant who knows the subject well or is documented in medical records
- * Meets the criteria for a diagnosis of probable Alzheimer's disease (AD) based on the National Institute of Neurological and Communicative Disorders - Stroke and the Alzheimer's Disease and Related Disorders Association (NINCDS-ADRDA) criteria for probable AD
- * Is receiving donepezil 10 mg daily for symptomatic treatment of cognitive impairment associated with AD. The dose level must be stable for at least 2 months prior to Screening. If receiving donepezil via a transdermal system (ie, patch), it should be a 10-mg/day dose and should switch prescription to a 10-mg oral daily dose, before enrollment
- * Has a reliable and competent trial partner/caregiver who has a close relationship with the participant, has face-to-face contact at least 3 days a week for a minimum of 6 waking hours a week, and is willing to accompany the participant, if desired, to study visits

4.2 Exclusion Criteria

- * History of clinically significant endocrine, gastrointestinal, cardiovascular, hematological, hepatic, immunological, renal, respiratory, genitourinary, or major neurological (including stroke and chronic seizures) abnormalities or diseases that are not under medical control over the past 2 months.
- * Has evidence of a clinically relevant or unstable psychiatric disorder, based on DSM-5 criteria, or has a history of clinically significant psychiatric disorder in the last 5 years. Generalized anxiety disorder, and/or insomnia under good control for ? 2 months on stable medical therapy may not be exclusionary.
- * History of cancer (malignancy). Participants with adequately treated disease deemed as "cured," or who, in the opinion of the study investigator, are highly unlikely to sustain a recurrence for the duration of the study, may be enrolled at the discretion of the investigator and Sponsor.
- * History of significant multiple and/or severe allergies (eg, food, drug, latex allergy), or has had an anaphylactic reaction or significant intolerability (ie, systemic allergic reaction) to prescription or nonprescription drugs or food.
- * Had a major surgery and/or donated or lost 1 unit of blood (approximately 500 mL) within 4 weeks prior to the prestudy (screening) visit.
- * Unable to refrain from or anticipates the use of any medication, including prescription and nonprescription drugs or herbal remedies beginning approximately 2 weeks (or 5 half-lives) prior to administration of the initial dose of study intervention, throughout the study, until the poststudy visit. There may be certain protocol-specified medications that are permitted.
- * The participant is a smoker and/or has used nicotine or nicotine-containing products (eg, nicotine patch and electronic cigarette) within 3 months of screening.
- * Consumes greater than 3 servings of alcoholic beverages per day. Participants who consume 4 servings of alcoholic beverages per day may be enrolled at the discretion of the investigator.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

* The participant is a regular user of cannabis, any illicit drugs or has a history of drug (including alcohol) abuse within approximately 2 years.

11. Study Identifiers & Governance

Primary ID:	1089974905016072149
Registry Number:	NCT06285240
Sponsor:	Merck
Study Title:	Efficacy and Safety of MK-1167 in Participants With Alzheimer's Disease Dementia Taking Stable Donepezil Treatment (MK-1167-007)

15. Sign-off & Approval

Principal Investigator:	_____	Date:	_____
Clinical Operations Lead:	_____	Date:	_____
Lead Statistician:	_____	Date:	_____