

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	China Faricimab Real World Evidence: Evaluation of Faricimab Effectiveness, Safety and Treatment Pattern, in Diabetic Macular Edema, Retinal Vein Occlusion and Neovascular Age-Related Macular Degeneration: The Farseeing Study
Short Title:	Real World Evidence in China: Faricimab Use in Diabetic Macular Edema, Retinal Vein Occlusion, and Neovascular Age-Related Macular Degeneration (The Farseeing Study)
Protocol Number:	415886411132371914
NCT Number:	NCT06439576
Sponsor Name:	Roche
Investigational Product:	faricimab
Phase:	Not Applicable
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Change in Visual Acuity from Baseline at Month 12: Visual acuity will be collected as measured per local practice and then will be converted to the approximate ETDRS letter score.
Secondary Obj.:	1. Visual Acuity Over Time 2. Change in Visual Acuity from Baseline Over Time 3. Central Subfield Thickness Over Time

2.2 Background & Rationale

The Farseeing Study will explore long-term effectiveness, safety, and treatment patterns among patients being treated with faricimab in real-world, routine clinical practice in China. It is a primary data collection, non-interventional, prospective and retrospective, multi-center study designed to collect real-world, long-term data to gain clinical evidence on faricimab, by observing cohorts of patients with neovascular age-related macular degeneration (nAMD), diabetic macular edema (DME), and retinal vein occlusion (RVO) who are receiving treatment with faricimab.

3. Study Design & Methodology

Design Framework:	Type: OBSERVATIONAL, Status: ACTIVE_NOT_RECRUITING
Target N:	1000

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

1. Have signed the informed consent
2. Female and male Chinese patients, who had been diagnosed with nAMD, DME, or RVO by CFP, OCT, FFA, ICGA, or OCTA

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

3. ≥50 years old for patients with nAMD, ≥18 years old both for patients with DME and RVO, at the time of signing informed consent (or the first administration of faricimab, whichever occurs first)
4. Patients for whom the decision to receive treatment with faricimab is made prior to and independent from study participation
5. Patients have received at least one faricimab treatment (the first dose) in the study eye

4.2 Exclusion Criteria

1. Patients not receiving treatment for nAMD/DME/RVO with faricimab according to the standard of care and in line with the current summary of product characteristics (SPC) / labeling in China
2. Active ocular inflammation or suspected / active ocular infection in either eye
3. Received any other anti-VEGF treatment after faricimab
4. Received any steroid treatment within 6 months (180 days) before the first faricimab treatment in the study eye
5. Any participation in any other clinical trials currently
6. Patients could not provide the clinical data (visual acuity and OCT images) within 2 weeks (14 days) before receiving the initial faricimab injection

11. Study Identifiers & Governance

Primary ID:	415886411132371914
Registry Number:	NCT06439576
Sponsor:	Roche
Study Title:	Real World Evidence in China: Faricimab Use in Diabetic Macular Edema, Retinal Vein Occlusion, and Neovascular Age-Related Macular Degeneration (The Farseeing Study)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____