

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	An Open-label, Multicenter Study to Evaluate Pharmacokinetics, Safety and Tolerability up to 6 Years of Intravenous Secukinumab Infusions in Pediatric Participants With Juvenile Psoriatic Arthritis
Short Title:	Study to Evaluate the Pharmacokinetics (PK), Safety and Tolerability up to 6 Years of Intravenous (i.v.) Secukinumab in Pediatric Participants With Juvenile Psoriatic Arthritis (JPsA).
Protocol Number:	41651310577497797
NCT Number:	NCT06751238
Sponsor Name:	Novartis
Investigational Product:	secukinumab
Phase:	PHASE1
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Maximum concentration on Day 1: Maximum concentration of secukinumab on Day 1 2. Maximum concentration at steady-state (C _{max} , ss): Maximum concentration at steady-state. 3. Minimum concentration at steady-state (C _{min} , ss): Minimum concentration at steady-state
Secondary Obj.:	1. Number of participants with Adverse Events (AEs) and Serious Adverse Events (SAEs) 2. Number of participants with clinically significant changes in clinical laboratory measures and vital signs.

2.2 Background & Rationale

The purpose of this study is to determine the PK, safety and tolerability of multiple doses of intravenous (i.v.) secukinumab in pediatric participants with JPsA

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE1, Status: RECRUITING
Target N:	20

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

* Participants parent's or legal representative(s) written informed consent and child's assent, if appropriate, must be obtained before any study related activity or assessment is performed. Of note, if the participant reaches age of consent (as per local law) during the study, they will also need to sign the corresponding study ICF (Informed Consent Form).

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Males and females ≥2 years old to <18 years old at the time of screening.
- * Confirmed diagnosis of JPsA according to the modified International League of Associations for Rheumatology (ILAR) classification criteria that must have occurred at least 6 months prior to screening.
- * Active JPsA disease defined as ≥3 active joints (swollen or if not swollen must be both tender and limited range of motion) at baseline (BSL).
- * Inadequate response (≥1 month) or intolerance to ≥1 Non-Steroidal Anti-Inflammatory Drug (NSAID) at screening.
- * Inadequate response (≥2 months) or intolerance to ≥1 Disease Modifying Anti-Rheumatic Drug (DMARD) at screening.
- * Concomitant use of the following second-line agents such as disease-modifying and/or immunosuppressive drugs to treat the JPsA will be allowed:

- * Stable dose of methotrexate (MTX) (maximum of 20 mg/ m² BSA/ week) for at least 4 weeks prior to the BSL visit, with folic/folinic acid supplementation (according to standard medical practice of the center).

- * Stable dose of an oral corticosteroid (CS) at a prednisone equivalent dose of <0.2 mg/kg/day or up to 10 mg/day maximum, whichever is less, for at least 7 days prior to BSL.

- * Stable dose of no more than one NSAID for at least 1 week prior to BSL.

Key Exclusion Criteria:

- * Participants with body weight less than 10 kg at screening.
- * Use of other investigational drugs within 4 weeks or 5 half-lives of BSL, or until the expected pharmacodynamic effect has returned to BSL, whichever is longer.
- * History of hypersensitivity to study drug or its excipients or to drugs of similar chemical classes.
- * Participants with active inflammatory bowel disease or active uveitis at screening or BSL.
- * Fulfilling diagnostic criteria for any International League of Associations for Rheumatology (ILAR) juvenile idiopathic arthritis (JIA) category other than JPsA at BSL.
- * Participants treated with prohibited medication
- * Participants taking any non-biologic DMARD at screening except for MTX.
- * Any medical or psychiatric condition which, in the investigator's opinion, would preclude the participant from adhering to the protocol or completing the study per protocol.

Other inclusion/exclusion criteria may apply

4.2 Exclusion Criteria

- * Participants with body weight less than 10 kg at screening.
- * Use of other investigational drugs within 4 weeks or 5 half-lives of BSL, or until the expected pharmacodynamic effect has returned to BSL, whichever is longer.
- * History of hypersensitivity to study drug or its excipients or to drugs of similar chemical classes.
- * Participants with active inflammatory bowel disease or active uveitis at screening or BSL.
- * Fulfilling diagnostic criteria for any International League of Associations for Rheumatology (ILAR) juvenile idiopathic arthritis (JIA) category other than JPsA at BSL.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Participants treated with prohibited medication
- * Participants taking any non-biologic DMARD at screening except for MTX.
- * Any medical or psychiatric condition which, in the investigator's opinion, would preclude the participant from adhering to the protocol or completing the study per protocol.

Other inclusion/exclusion criteria may apply

11. Study Identifiers & Governance

Primary ID: 41651310577497797

Registry Number: NCT06751238

Sponsor: Novartis

Study Title: Study to Evaluate the Pharmacokinetics (PK), Safety and Tolerability up to 6 Years of Intravenous (i.v.) Secukinumab in Pediatric Participants With Juvenile Psoriatic Arthritis (JPsA).

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____